



SNX3 Polyclonal Antibody

Catalog No	BYab-00733
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB;ELISA;IHC
Gene Name	SNX3
Protein Name	Sorting nexin-3
Immunogen	The antiserum was produced against synthesized peptide derived from human SNX3. AA range:91-140
Specificity	SNX3 Polyclonal Antibody detects endogenous levels of SNX3 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000;IHC-p 1:50-300; ELISA 2000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	SNX3; Sorting nexin-3; Protein SDP3
Observed Band	18kD
Cell Pathway	Early endosome . Cytoplasmic vesicle, phagosome . Colocalizes to clathrin-coated endosomal vesicles morphologically distinct from retromer-decorated non-branched endosomal tubule structures (PubMed:21725319) Colocalizes with EEA1 on nascent phagosomes in dendritic cells but competes with EEA1 for binding to phagosomal membrane (PubMed:23237080). In the case of Salmonella enterica infection localizes to Salmonella-containing vacuoles (SCVs) from which SNX3-containing tubules form 30-60 min after infection (PubMed:20482551). .
Tissue Specificity	Brain,Colon,Epithelium,Pancreas,Platelet,Skin,
Function	disease:A chromosomal aberration disrupting SNX3 may be a cause of microphthalmia syndromic type 8 (MCOPS8) [MIM:601349]. Translocation t(6;13)(q21;q12). Microphthalmia is a clinically heterogeneous disorder of eye formation, ranging from small size of a single eye to complete bilateral absence of ocular tissues (anophthalmia). In many cases, microphthalmia/anophthalmia

Nanjing BYabscience technology Co.,Ltd



occurs in association with syndromes that include non-ocular abnormalities. MCOPS8 is a very rare congenital syndrome characterized by microcephaly, microphthalmia, ectrodactyly of the lower limbs and prognathism. Intellectual deficit has been reported. function: May be involved in several stages of intracellular trafficking. similarity: Belongs to the sorting nexin family. similarity: Contains 1 PX (phox homology) domain.

Background

This gene encodes a member of the sorting nexin family. Members of this family contain a phox (PX) domain, which is a phosphoinositide binding domain, and are involved in intracellular trafficking. This protein does not contain a coiled coil region, like most family members. This protein interacts with phosphatidylinositol-3-phosphate, and is involved in protein trafficking. A pseudogene of this gene is present on the sex chromosomes. Alternative splicing results in multiple transcript variants encoding distinct isoforms. [provided by RefSeq, Jul 2014],

matters needing attention


Avoid repeated freezing and thawing!

Usage suggestions


This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.




Products Images



Western blot analysis of SNX3 Antibody. The lane on the right is blocked with the SNX3 peptide.



Western blot analysis of the lysates from HUVEC cells using SNX3 antibody.



Immunohistochemical analysis of paraffin-embedded human tonsil. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).

Nanjing BYabscience technology Co.,Ltd