



Socius Polyclonal Antibody

Catalog No	BYab-00735
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB;IHC;IF;ELISA
Gene Name	UBXN11
Protein Name	UBX domain-containing protein 11
Immunogen	The antiserum was produced against synthesized peptide derived from human UBXD5. AA range:131-180
Specificity	Socius Polyclonal Antibody detects endogenous levels of Socius protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/20000. Not yet tested in other applications.
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	UBXN11; SOC; UBXD5; PP2243; UBX domain-containing protein 11; Colorectal tumor-associated antigen COA-1; Socius; UBX domain-containing protein 5
Observed Band	57kD
Cell Pathway	Cytoplasm, cytoskeleton .
Tissue Specificity	Colon cancer,Small intestine,Synovium,Testis,
Function	function:May be involved in the reorganization of actin cytoskeleton mediated by RND1, RND2 AND RND3. Promotes RHOA activation mediated by GNA12 and GNA13.,polymorphism:The number of repeats is polymorphic and varies from 1 to 4.,similarity:Contains 1 SEP domain.,similarity:Contains 1 UBX domain.,subunit:Interacts with GNA12, GNA13, RND1, RND2 AND RND3.,
Background	This gene encodes a protein with a divergent C-terminal UBX domain. The homologous protein in the rat interacts with members of the Rnd subfamily of Rho GTPases at the cell periphery through its C-terminal region. It also interacts with

Nanjing BYabscience technology Co.,Ltd



several heterotrimeric G proteins through their G-alpha subunits and promotes Rho GTPase activation. It is proposed to serve a bidirectional role in the promotion and inhibition of Rho activity through upstream signaling pathways. The 3' coding sequence of this gene contains a polymorphic region of 24 nt tandem repeats. Several transcripts containing between 1.5 and five repeat units have been reported. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jul 2008],

matters needing attention


Avoid repeated freezing and thawing!

Usage suggestions


This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.



Products Images



Immunohistochemistry analysis of paraffin-embedded human lung carcinoma tissue, using UBXD5 Antibody. The picture on the right is blocked with the synthesized peptide.



Western blot analysis of lysates from HepG2 cells, using UBXD5 Antibody. The lane on the right is blocked with the synthesized peptide.