



TRAPPC5 Polyclonal Antibody

Catalog No BYab-00751 Isotype IgG Reactivity Human;Mouse;Rat Applications IHC;IF;ELISA Gene Name TRAPPC5 Protein Name Trafficking protein particle complex subunit 5 Immunogen Synthesized peptide derived from the Internal region of human TRAPPC5. Specificity TRAPPC5 Polyclonal Antibody detects endogenous levels of TRAPPC5 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution IHC: 1/100 - 1/300. ELISA: 1/5000 IF 1:50-200 Concentration 1 mg/ml Purity ≥90% Storage Stability -20*C/1 year Synonyms TRAPPC5; Trafficking protein particle complex subunit 5 Observed Band Cell Pathway Golgi apparatus, cis-Golgi network . Endoplasmic reticulum . Tissue Specificity Skin, Function function:May play a role in vesicular transport from endoplasmic reticulum to Golgi, similarity. Belongs to the TRAPP small subunits family. BET3 sub		
Reactivity Human;Mouse;Rat Applications IHC;IF;ELISA Gene Name TRAPPC5 Protein Name Trafficking protein particle complex subunit 5 Immunogen Synthesized peptide derived from the Internal region of human TRAPPC5. Specificity TRAPPC5 Polyclonal Antibody detects endogenous levels of TRAPPC5 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution IHC: 1/100 - 1/300. ELISA: 1/5000 IF 1:50-200 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms TRAPPC5; Trafficking protein particle complex subunit 5 Observed Band Cell Pathway Golgi apparatus, cis-Golgi network . Endoplasmic reticulum . Tissue Specificity Skin, Function function:May play a role in vesicular transport from endoplasmic reticulum to Golgi, similarity:Belongs to the TRAPP small subunits family. BET3 subfamily. subunit.Part of the multisubunit TRAPP (transport protein particle) complex., Background Avoid repeated freezing and thawing!	Catalog No	BYab-00751
Applications IHC;IF;ELISA Gene Name TRAPPC5 Protein Name Trafficking protein particle complex subunit 5 Immunogen Synthesized peptide derived from the Internal region of human TRAPPC5. Specificity TRAPPC5 Polyclonal Antibody detects endogenous levels of TRAPPC5 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Puriffication The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution IHC: 1/100 - 1/300. ELISA: 1/5000 IF 1:50-200 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms TRAPPC5; Trafficking protein particle complex subunit 5 Observed Band Cell Pathway Golgi apparatus, cis-Golgi network . Endoplasmic reticulum . Tissue Specificity Skin, Function function:May play a role in vesicular transport from endoplasmic reticulum to Golgi, similarity:Belongs to the TRAPP small subunits family. BET3 subfamily. subunit.Part of the multisubunit TRAPP (transport protein particle) complex., Background function:May play a role in vesicular transport from endoplasmic reticulum to Golgi, similarity:Belongs to the TRAPP small subunits family. BET3 subfamily., subunit.Part of the multisubunit TRAPP (transport protein particle) complex., Matters needing Avoid repeated freezing and thawing!	Isotype	IgG
Gene Name TRAPPC5 Protein Name Trafficking protein particle complex subunit 5 Immunogen Synthesized peptide derived from the Internal region of human TRAPPC5. Specificity TRAPPC5 Polyclonal Antibody detects endogenous levels of TRAPPC5 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution IHC: 1/100 - 1/300. ELISA: 1/5000 IF 1:50-200 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms TRAPPC5; Trafficking protein particle complex subunit 5 Observed Band Cell Pathway Golgi apparatus, cis-Golgi network . Endoplasmic reticulum . Tissue Specificity Skin, Function function:May play a role in vesicular transport from endoplasmic reticulum to Golgi, similarity:Belongs to the TRAPP small subunits family. BET3 subfamily, subunit:Part of the multisubunit TRAPP (transport protein particle) complex., Background function:May play a role in vesicular transport from endoplasmic reticulum to Golgi, similarity:Belongs to the TRAPP small subunits family. BET3 sub	Reactivity	Human;Mouse;Rat
Protein Name Trafficking protein particle complex subunit 5 Immunogen Synthesized peptide derived from the Internal region of human TRAPPC5. Specificity TRAPPC5 Polyclonal Antibody detects endogenous levels of TRAPPC5 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution IHC: 1/100 - 1/300. ELISA: 1/5000 IF 1:50-200 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms TRAPPC5; Trafficking protein particle complex subunit 5 Observed Band Cell Pathway Golgi apparatus, cis-Golgi network . Endoplasmic reticulum . Tissue Specificity Skin, Function function:May play a role in vesicular transport from endoplasmic reticulum to Golgisimilarity:Belongs to the TRAPP small subunits family. BET3 subfamily., subunit:Part of the multisubunit TRAPP (transport protein particle) complex., matters needing Avoid repeated freezing and thawing!	Applications	IHC;IF;ELISA
Immunogen Synthesized peptide derived from the Internal region of human TRAPPC5. Specificity TRAPPC5 Polyclonal Antibody detects endogenous levels of TRAPPC5 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution IHC: 1/100 - 1/300. ELISA: 1/5000 IF 1:50-200 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms TRAPPC5; Trafficking protein particle complex subunit 5 Observed Band Cell Pathway Golgi apparatus, cis-Golgi network . Endoplasmic reticulum . Tissue Specificity Skin, Function function:May play a role in vesicular transport from endoplasmic reticulum to Golgi, similarity:Belongs to the TRAPP small subunits family. BET3 subfamily, subunit:Part of the multisubunit TRAPP (transport protein particle) complex., matters needing Avoid repeated freezing and thawing!	Gene Name	TRAPPC5
Specificity TRAPPC5 Polyclonal Antibody detects endogenous levels of TRAPPC5 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution IHC: 1/100 - 1/300. ELISA: 1/5000 IF 1:50-200 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms TRAPPC5; Trafficking protein particle complex subunit 5 Observed Band Cell Pathway Gelgi apparatus, cis-Gelgi network . Endoplasmic reticulum . Tissue Specificity Skin, Function function:May play a role in vesicular transport from endoplasmic reticulum to Golgi., similarity. Belongs to the TRAPP small subunits family. BET3 subfamily., subunit:Part of the multisubunit TRAPP (transport protein particle) complex., subunit:Part of the multisubunit TRAPP (transport protein particle) complex. matters needing Avoid repeated freezing and thawing!	Protein Name	Trafficking protein particle complex subunit 5
Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution IHC: 1/100 - 1/300. ELISA: 1/5000 IF 1:50-200 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms TRAPPC5; Trafficking protein particle complex subunit 5 Observed Band Cell Pathway Golgi apparatus, cis-Golgi network . Endoplasmic reticulum . Tissue Specificity Skin, Function Golgi, similarity: Belongs to the TRAPP small subunits family. BET3 subfamily, subunit: Part of the multisubunit TRAPP (transport protein particle) complex., Background function:May play a role in vesicular transport from endoplasmic reticulum to Golgi, similarity: Belongs to the TRAPP small subunits family. BET3 subfamily, subunit: Part of the multisubunit TRAPP (transport protein particle) complex., matters needing Avoid repeated freezing and thawing!	Immunogen	Synthesized peptide derived from the Internal region of human TRAPPC5.
Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution IHC: 1/100 - 1/300. ELISA: 1/5000 IF 1:50-200 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms TRAPPC5; Trafficking protein particle complex subunit 5 Observed Band Cell Pathway Golgi apparatus, cis-Golgi network . Endoplasmic reticulum . Tissue Specificity Skin, Function function:May play a role in vesicular transport from endoplasmic reticulum to Golgi, similarity:Belongs to the TRAPP small subunits family. BET3 subfamily., subunit:Part of the multisubunit TRAPP (transport protein particle) complex., subfamily., subunit:Part of the multisubunit TRAPP (transport protein particle) complex., aubtamily., subunit:Part of the multisubunit TRAPP (transport protein particle) complex. matters needing Avoid repeated freezing and thawing!	Specificity	TRAPPC5 Polyclonal Antibody detects endogenous levels of TRAPPC5 protein.
Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution IHC: 1/100 - 1/300. ELISA: 1/5000 IF 1:50-200 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms TRAPPC5; Trafficking protein particle complex subunit 5 Observed Band Cell Pathway Golgi apparatus, cis-Golgi network . Endoplasmic reticulum . Tissue Specificity Skin, Function function:May play a role in vesicular transport from endoplasmic reticulum to Golgi., similarity:Belongs to the TRAPP small subunits family. BET3 subfamily., subunit:Part of the multisubunit TRAPP (transport protein particle) complex., function:May play a role in vesicular transport from endoplasmic reticulum to Golgi., similarity:Belongs to the TRAPP small subunits family. BET3 subfamily., subunit:Part of the multisubunit TRAPP (transport protein particle) complex., Matters needing Avoid repeated freezing and thawing!	Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen. Dilution IHC: 1/100 - 1/300. ELISA: 1/5000 IF 1:50-200 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms TRAPPC5; Trafficking protein particle complex subunit 5 Observed Band Cell Pathway Golgi apparatus, cis-Golgi network . Endoplasmic reticulum . Tissue Specificity Skin, Function function:May play a role in vesicular transport from endoplasmic reticulum to Golgi.,similarity:Belongs to the TRAPP small subunits family. BET3 subfamily.,subunit:Part of the multisubunit TRAPP (transport protein particle) complex., Background function:May play a role in vesicular transport from endoplasmic reticulum to Golgi,similarity:Belongs to the TRAPP small subunits family. BET3 subfamily.,subunit:Part of the multisubunit TRAPP (transport protein particle) complex., Matters needing Avoid repeated freezing and thawing!	Source	Polyclonal, Rabbit,IgG
Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms TRAPPC5; Trafficking protein particle complex subunit 5 Observed Band Cell Pathway Golgi apparatus, cis-Golgi network . Endoplasmic reticulum . Tissue Specificity Skin, Function function:May play a role in vesicular transport from endoplasmic reticulum to Golgi.,similarity:Belongs to the TRAPP small subunits family. BET3 subfamily.,subunit:Part of the multisubunit TRAPP (transport protein particle) complex., Background function:May play a role in vesicular transport from endoplasmic reticulum to Golgi.,similarity:Belongs to the TRAPP small subunits family. BET3 subfamily. Betall subunits family. BET3 subfamily. subunit:Part of the multisubunit TRAPP (transport protein particle) complex., matters needing Avoid repeated freezing and thawing!	Purification	
Purity ≥90% Storage Stability -20°C/1 year Synonyms TRAPPC5; Trafficking protein particle complex subunit 5 Observed Band Cell Pathway Golgi apparatus, cis-Golgi network . Endoplasmic reticulum . Tissue Specificity Skin, Function function:May play a role in vesicular transport from endoplasmic reticulum to Golgi.,similarity:Belongs to the TRAPP small subunits family. BET3 subfamily.,subunit:Part of the multisubunit TRAPP (transport protein particle) complex., Background function:May play a role in vesicular transport from endoplasmic reticulum to Golgi.,similarity:Belongs to the TRAPP small subunits family. BET3 subfamily.,subunit:Part of the multisubunit TRAPP (transport protein particle) complex., matters needing Avoid repeated freezing and thawing!	Dilution	IHC: 1/100 - 1/300. ELISA: 1/5000 IF 1:50-200
Synonyms TRAPPC5; Trafficking protein particle complex subunit 5 Observed Band Cell Pathway Golgi apparatus, cis-Golgi network . Endoplasmic reticulum . Tissue Specificity Skin, Function function:May play a role in vesicular transport from endoplasmic reticulum to Golgi.,similarity:Belongs to the TRAPP small subunits family. BET3 subfamily.,subunit:Part of the multisubunit TRAPP (transport protein particle) complex., Background function:May play a role in vesicular transport from endoplasmic reticulum to Golgi.,similarity:Belongs to the TRAPP small subunits family. BET3 subfamily.,subunit:Part of the multisubunit TRAPP (transport protein particle) complex., matters needing Avoid repeated freezing and thawing!	Concentration	1 mg/ml
Synonyms TRAPPC5; Trafficking protein particle complex subunit 5 Observed Band Cell Pathway Golgi apparatus, cis-Golgi network . Endoplasmic reticulum . Tissue Specificity Skin, Function function:May play a role in vesicular transport from endoplasmic reticulum to Golgi.,similarity:Belongs to the TRAPP small subunits family. BET3 subfamily.,subunit:Part of the multisubunit TRAPP (transport protein particle) complex., Background function:May play a role in vesicular transport from endoplasmic reticulum to Golgi.,similarity:Belongs to the TRAPP small subunits family. BET3 subfamily.,subunit:Part of the multisubunit TRAPP (transport protein particle) complex., matters needing Avoid repeated freezing and thawing!	Purity	≥90%
Observed Band Cell Pathway Golgi apparatus, cis-Golgi network . Endoplasmic reticulum . Tissue Specificity Skin, Function function:May play a role in vesicular transport from endoplasmic reticulum to Golgi.,similarity:Belongs to the TRAPP small subunits family. BET3 subfamily.,subunit:Part of the multisubunit TRAPP (transport protein particle) complex., Background function:May play a role in vesicular transport from endoplasmic reticulum to Golgi.,similarity:Belongs to the TRAPP small subunits family. BET3 subfamily.,subunit:Part of the multisubunit TRAPP (transport protein particle) complex., matters needing Avoid repeated freezing and thawing!	Storage Stability	-20°C/1 year
Cell Pathway Golgi apparatus, cis-Golgi network . Endoplasmic reticulum . Skin, Function function:May play a role in vesicular transport from endoplasmic reticulum to Golgi.,similarity:Belongs to the TRAPP small subunits family. BET3 subfamily.,subunit:Part of the multisubunit TRAPP (transport protein particle) complex., Background function:May play a role in vesicular transport from endoplasmic reticulum to Golgi.,similarity:Belongs to the TRAPP small subunits family. BET3 subfamily.,subunit:Part of the multisubunit TRAPP (transport protein particle) complex., matters needing Avoid repeated freezing and thawing!	Synonyms	TRAPPC5; Trafficking protein particle complex subunit 5
Tissue Specificity Skin, Function function:May play a role in vesicular transport from endoplasmic reticulum to Golgi.,similarity:Belongs to the TRAPP small subunits family. BET3 subfamily.,subunit:Part of the multisubunit TRAPP (transport protein particle) complex., Background function:May play a role in vesicular transport from endoplasmic reticulum to Golgi.,similarity:Belongs to the TRAPP small subunits family. BET3 subfamily.,subunit:Part of the multisubunit TRAPP (transport protein particle) complex., matters needing Avoid repeated freezing and thawing!	Observed Band	
Function function:May play a role in vesicular transport from endoplasmic reticulum to Golgi.,similarity:Belongs to the TRAPP small subunits family. BET3 subfamily.,subunit:Part of the multisubunit TRAPP (transport protein particle) complex., function:May play a role in vesicular transport from endoplasmic reticulum to Golgi.,similarity:Belongs to the TRAPP small subunits family. BET3 subfamily.,subunit:Part of the multisubunit TRAPP (transport protein particle) complex., matters needing Avoid repeated freezing and thawing!	Cell Pathway	Golgi apparatus, cis-Golgi network . Endoplasmic reticulum .
Golgi.,similarity:Belongs to the TRAPP small subunits family. BET3 subfamily.,subunit:Part of the multisubunit TRAPP (transport protein particle) complex., Background function:May play a role in vesicular transport from endoplasmic reticulum to Golgi.,similarity:Belongs to the TRAPP small subunits family. BET3 subfamily.,subunit:Part of the multisubunit TRAPP (transport protein particle) complex., matters needing Avoid repeated freezing and thawing!	Tissue Specificity	Skin,
Golgi.,similarity:Belongs to the TRAPP small subunits family. BET3 subfamily.,subunit:Part of the multisubunit TRAPP (transport protein particle) complex., matters needing Avoid repeated freezing and thawing!	Function	Golgi.,similarity:Belongs to the TRAPP small subunits family. BET3 subfamily.,subunit:Part of the multisubunit TRAPP (transport protein particle)
	Background	Golgi.,similarity:Belongs to the TRAPP small subunits family. BET3 subfamily.,subunit:Part of the multisubunit TRAPP (transport protein particle)
	matters needing attention	Avoid repeated freezing and thawing!

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658




国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务, 欢迎咨询



Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Immunohistochemical analysis of paraffin-embedded human Gastric adenocarcinoma. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).

Nanjing BYabscience technology Co.,Ltd