



Histone H2A.Z (Acetyl Lys12) rabbit pAb

Catalog No	BYab-00887
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB; ELISA
Gene Name	H2AFZ H2AZ
Protein Name	Histone H2A.Z (Acetyl Lys12)
Immunogen	Synthesized peptide derived from human Histone H2A.Z (Acetyl Lys12)
Specificity	This antibody detects endogenous levels of Human,Mouse,Rat Histone H2A.Z (Acetyl Lys12)
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1:1000-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	Histone H2A.Z (H2A/z)
Observed Band	15kD
Cell Pathway	Nucleus. Chromosome.
Tissue Specificity	
Function	DNA packaging, chromatin organization, chromatin assembly or disassembly, nucleosome assembly, chromatin assembly, cellular macromolecular complex subunit organization, cellular macromolecular complex assembly,nucleosome organization, macromolecular complex subunit organization, chromosome organization, macromolecular complex assembly, protein-DNA complex assembly,
Background	function:Variant histone H2A which replaces conventional H2A in a subset of nucleosomes. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a

Nanjing BYabscience technology Co.,Ltd



complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling. May be involved in the formation of constitutive heterochromatin. May be required for chromosome segregation during cell division.,mass spectrometry:Monoisotopic, not modified PubMed:16457589,PTM:Acetylated on Lys-5, Lys-8 and Lys-12 during interphase. Acetylation disappears at mitosis.,PTM:Monoubiquitination of Lys-122 gives a specific tag for epigenetic transcriptional repression.,PTM:Not phosphorylated.,similarity:Belongs to the histone H2A family.,subunit:The nucleosome is a histone octamer containing two molecules each of H2A, H2B, H3 and H4 assembled in one H3-H4 heterotetramer and two H2A-H2B heterodimers. The octamer wraps approximately 147 bp of DNA. H2A or its variant H2AFZ forms an heterodimer with H2B. H2AFZ interacts with INCENP.,

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images