



ESET Monoclonal Antibody

Catalog No	BYab-00973
Isotype	IgG
Reactivity	Human;Mouse;Monkey
Applications	WB;IF;ELISA
Gene Name	SETDB1
Protein Name	Histone-lysine N-methyltransferase SETDB1
Immunogen	Purified recombinant fragment of human ESET expressed in E. Coli.
Specificity	ESET Monoclonal Antibody detects endogenous levels of ESET protein.
Formulation	Ascitic fluid containing 0.03% sodium azide,0.5% BSA, 50%glycerol.
Source	Monoclonal, Mouse
Purification	Affinity purification
Dilution	Western Blot: 1/500 - 1/2000. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/10000. Not yet tested in other applications.
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	SETDB1; KIAA0067; KMT1E; Histone-lysine N-methyltransferase SETDB1; ERG-associated protein with SET domain; ESET; Histone H3-K9 methyltransferase 4; H3-K9-HMTase 4; Lysine N-methyltransferase 1E; SET domain bifurcated 1
Observed Band	
Cell Pathway	Nucleus . Cytoplasm . Chromosome. Associated with non-pericentromeric regions of chromatin. Excluded from nucleoli and islands of condensed chromatin. .
Tissue Specificity	Widely expressed. High expression in testis.
Function	catalytic activity:S-adenosyl-L-methionine + histone L-lysine = S-adenosyl-L-homocysteine + histone N(6)-methyl-L-lysine.,domain:The pre-SET, SET and post-SET domains are all required for methyltransferase activity. The 347-amino-acid insertion in the SET domain has no effect on the catalytic activity.,function:Histone methyltransferase that specifically trimethylates 'Lys-9' of histone H3. H3 'Lys-9' trimethylation represents a specific tag for epigenetic transcriptional repression by recruiting HP1 (CBX1, CBX3 and/or CBX5) proteins to methylated histones. Mainly functions in euchromatin regions, thereby playing a central role in the silencing of euchromatic genes. H3 'Lys-9'

Nanjing BYabscience technology Co.,Ltd



trimethylation is coordinated with DNA methylation. Probably forms a complex with MBD1 and ATF7IP that represses transcription and couples DNA methylation and histone 'Lys-9' trimethylation. Its activity is depende

Background

SET domain bifurcated 1(SETDB1) Homo sapiens This gene encodes a histone methyltransferase which regulates histone methylation, gene silencing, and transcriptional repression. This gene has been identified as a target for treatment in Huntington Disease, given that gene silencing and transcription dysfunction likely play a role in the disease pathogenesis. Alternatively spliced transcript variants of this gene have been described.[provided by RefSeq, Jun 2011],

matters needing attention


Avoid repeated freezing and thawing!

Usage suggestions


This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.



Products Images



Western Blot analysis using ESET Monoclonal Antibody against MCF-7 (1), T47D (2), HEK293 (3), JURKAT (4), NIH/3T3 (5), F9 (6), RAW246.7 (7) and Cos7 (8) cell lysate.



Immunofluorescence analysis of LOVO cells using ESET Monoclonal Antibody (green). Red: Actin filaments have been labeled with Alexa Fluor-555 phalloidin.