



# CHD3(C-term) mouse mAb

<b>Catalog No</b>	BYab-01074
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human
<b>Applications</b>	WB;IHC;FC
<b>Gene Name</b>	chd3
<b>Protein Name</b>	
<b>Immunogen</b>	Purified recombinant human CHD3 (C-terminus) protein fragments expressed in E.coli.
<b>Specificity</b>	This antibody detects endogenous levels of CHD3 (C-terminus) and does not cross-react with related proteins.
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source</b>	Monoclonal, Mouse
<b>Purification</b>	The antibody was affinity-purified from mouse ascites by affinity-chromatography using epitope-specific immunogen.
<b>Dilution</b>	wb 1:1000 1:500-1:1000 fcm 1:100
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	226;2600010P09Rik;AF020312;ATP dependent helicase CHD3;ATP-dependent helicase Chd3;CHD-3;Chd3;CHD3_HUMAN;Chd7;Chromodomain-helicase-DNA-binding protein 3;hZFH;MGC40857;Mi 2 autoantigen 240 kDa protein;Mi 2a;Mi-2 autoantigen 240 kDa protein;Mi2 ALPHA;Mi2-alpha; Prp7;Prp9 1;ZFH;Zinc finger helicase;zinc-finger helicase (Snf2-like).
<b>Observed Band</b>	260kD
<b>Cell Pathway</b>	Nucleus, PML body . Nucleus . Cytoplasm, cytoskeleton, microtubule organizing center, centrosome . Associates with centrosomes in interphase and mitosis. .
<b>Tissue Specificity</b>	Widely expressed.
<b>Function</b>	disease:One of the main antigens reacting with anti-MI-2 positive sera of dermatomyositis.,function:Probable transcription regulator.,sequence caution:Differs from position 1967 onward for unknown reasons.,similarity:Belongs to the SNF2/RAD54 helicase

Nanjing BYabscience technology Co.,Ltd



family.,similarity:Contains 1 helicase ATP-binding domain.,similarity:Contains 1 helicase C-terminal domain.,similarity:Contains 2 chromo domains.,similarity:Contains 2 PHD-type zinc fingers.,subunit:Central component of the nucleosome remodeling and histone deacetylase (NuRD) repressive complex. Interacts with TRIM28 and SERBP1. Interacts via its C-terminal region with HBP4.,tissue specificity:Widely expressed.,

**Background**

This gene encodes a member of the CHD family of proteins which are characterized by the presence of chromo (chromatin organization modifier) domains and SNF2-related helicase/ATPase domains. This protein is one of the components of a histone deacetylase complex referred to as the Mi-2/NuRD complex which participates in the remodeling of chromatin by deacetylating histones. Chromatin remodeling is essential for many processes including transcription. Autoantibodies against this protein are found in a subset of patients with dermatomyositis. Three alternatively spliced transcripts encoding different isoforms have been described. [provided by RefSeq, Jul 2008],

**matters needing attention**


Avoid repeated freezing and thawing!

**Usage suggestions**


This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.




## Products Images




Western blot detection of CHD3 (C-terminus) in Y79 and Raji cell lysates using CHD3 (C-terminus) mouse mAb (1:1000 diluted). Predicted band size: 226KDa. Observed band size: 260KDa.



IHC of paraffin-embedded human colon using anti-CHD3 (C-terminus) diluted 1/500-1/1000.



Flow Cytometry analysis of K562 cells stained with CHD3 (red, 1/100 dilution), followed by FITC-conjugated goat anti-mouse IgG. Blue line histogram represents the isotype control, normal mouse IgG.



Immunohistochemical analysis of paraffin-embedded human tonsil. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).