



SMAD5(C-term) mouse mAb

Catalog No	BYab-01077
Isotype	IgG
Reactivity	Human
Applications	WB;ICC;FC
Gene Name	smad5
Protein Name	
Immunogen	Purified recombinant human SMAD5 (C-terminus) protein fragments expressed in E.coli.
Specificity	This antibody detects endogenous levels of SMAD5 (C-terminus) and does not cross-react with related proteins.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse
Purification	The antibody was affinity-purified from mouse ascites by affinity-chromatography using epitope-specific immunogen.
Dilution	wb 1:1000 icc 1:75 fcm 1:100
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	DKFZp781C1895;DKFZp781O1323;Dwfc;hSmad 5;hSmad5;JV5 1;JV5-1;MAD homolog 5;MAD mothers against decapentaplegic homolog 5;MAD, mothers against decapentaplegic homolog 5;MADH 5;MADH5;Mothers against decapentaplegic homolog 5;Mothers against DPP homolog 5;MusMLP; SMA and MAD related protein 5;SMAD 5;SMAD family member 5;SMAD mothers against DPP homolog 5;Smad5;Smad5;SMAD5_HUMAN.
Observed Band	60kD
Cell Pathway	Cytoplasm. Nucleus. Cytoplasmic in the absence of ligand. Migrates to the nucleus when complexed with SMAD4.
Tissue Specificity	Ubiquitous.
Function	function:Transcriptional modulator activated by BMP (bone morphogenetic proteins) type 1 receptor kinase. SMAD5 is a receptor-regulated SMAD (R-SMAD).,PTM:Phosphorylated on serine by BMP (bone morphogenetic proteins) type 1 receptor kinase.,PTM:Ubiquitin-mediated proteolysis by

Nanjing BYabscience technology Co.,Ltd



SMAD-specific E3 ubiquitin ligase SMURF1.,similarity:Belongs to the dwarfin/SMAD family.,similarity:Contains 1 MH1 (MAD homology 1) domain.,similarity:Contains 1 MH2 (MAD homology 2) domain.,subcellular location:Cytoplasmic in the absence of ligand. Migrates to the nucleus when complexed with SMAD4.,subunit:May form trimers with the co-SMAD SMAD4. Interacts with PEBP2-alpha subunit and SMURF1. Interacts with SUV39H1 and SUV39H2.,tissue specificity:Ubiquitous.,

Background

The protein encoded by this gene is involved in the transforming growth factor beta signaling pathway that results in an inhibition of the proliferation of hematopoietic progenitor cells. The encoded protein is activated by bone morphogenetic proteins type 1 receptor kinase, and may be involved in cancer. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Feb 2014],

matters needing attention


Avoid repeated freezing and thawing!

Usage suggestions


This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.



Products Images




Western blot detection of SMAD5 (C-terminus) in HeLa, Jurkat and K562 cell lysates using SMAD5 (C-terminus) mouse mAb (1:1000 diluted). Predicted band size: 52KDa. Observed band size: 60KDa.



Immunocytochemistry of HeLa cells using anti-SMAD5 (C-terminus) mouse mAb diluted 1:75.



Flow Cytometry analysis of Jurkat cells stained with SMAD5 (red, 1/100 dilution), followed by FITC-conjugated goat anti-mouse IgG. Black line histogram represents the isotype control, normal mouse IgG.



Nanjing BYabscience technology Co.,Ltd