




JunB (phospho-Thr102/Thr104) rabbit pAb

Catalog No	BYab-01455
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB;IHC
Gene Name	JUNB
Protein Name	JunB (Thr102/Thr104)
Immunogen	Synthesized phosho peptide around human JunB (Thr102 and Thr104)
Specificity	This antibody detects endogenous levels of Human JunB (phospho-Thr102 or Thr104)
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1:500-2000;IHC-p 1:50-300
Concentration	1 mg/ml
-	
Purity	≥90%
	≥90% -20°C/1 year
Purity	
Purity Storage Stability	-20°C/1 year
Purity Storage Stability Synonyms	-20°C/1 year Transcription factor jun-B
Purity Storage Stability Synonyms Observed Band	-20°C/1 year Transcription factor jun-B 36kD
Purity Storage Stability Synonyms Observed Band Cell Pathway	-20°C/1 year Transcription factor jun-B 36kD Nucleus.
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity	-20°C/1 year Transcription factor jun-B 36kD Nucleus. Adipose tissue,Epithelium,Pancreas,Placenta,Prostat function:Transcription factor involved in regulating gene activity following the primary growth factor response. Binds to the DNA sequence 5'-TGA[CG]TCA-3'.,induction:By growth factors.,similarity:Belongs to the bZIP family.,similarity:Belongs to the bZIP family. Jun subfamily.,similarity:Contains 1 bZIP domain.,subunit:Binds DNA as an homodimer or as an heterodimer with

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658









Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western blot analysis of lysates from K562 cells, primary antibody was diluted at 1:1000, 4° over night



Immunohistochemical analysis of paraffin-embedded human Gastric adenocarcinoma. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658