



ASCL1 Polyclonal Antibody

han;Mouse;Rat IHC;IF;ELISA CL1 aete-scute homolog 1 antiserum was produced against synthesized peptide derived from human CL1. AA range:109-158 CL1 Polyclonal Antibody detects endogenous levels of ASCL1 protein. id in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. /clonal, Rabbit,IgG antibody was affinity-purified from rabbit antiserum by ity-chromatography using epitope-specific immunogen.
HC;IF;ELISA CL1 aete-scute homolog 1 antiserum was produced against synthesized peptide derived from human CL1. AA range:109-158 CL1 Polyclonal Antibody detects endogenous levels of ASCL1 protein. id in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. /clonal, Rabbit,IgG antibody was affinity-purified from rabbit antiserum by
CL1 aete-scute homolog 1 antiserum was produced against synthesized peptide derived from human CL1. AA range:109-158 CL1 Polyclonal Antibody detects endogenous levels of ASCL1 protein. iid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. /clonal, Rabbit,IgG antibody was affinity-purified from rabbit antiserum by
aete-scute homolog 1 antiserum was produced against synthesized peptide derived from human CL1. AA range:109-158 CL1 Polyclonal Antibody detects endogenous levels of ASCL1 protein. iid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. /clonal, Rabbit,IgG antibody was affinity-purified from rabbit antiserum by
antiserum was produced against synthesized peptide derived from human CL1. AA range:109-158 CL1 Polyclonal Antibody detects endogenous levels of ASCL1 protein. iid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. //clonal, Rabbit,IgG antibody was affinity-purified from rabbit antiserum by
CL1. AA range:109-158 CL1 Polyclonal Antibody detects endogenous levels of ASCL1 protein. iid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. /clonal, Rabbit,IgG antibody was affinity-purified from rabbit antiserum by
antibody was affinity-purified from rabbit antiserum by
clonal, Rabbit,IgG antibody was affinity-purified from rabbit antiserum by
antibody was affinity-purified from rabbit antiserum by
: 1/500 - 1/2000. IHC: 1/100 - 1/300. ELISA: 1/40000 IF 1:50-200
g/ml
%
C/1 year
CL1; ASH1; BHLHA46; HASH1; Achaete-scute homolog 1; ASH-1; hASH1; ss A basic helix-loop-helix protein 46; bHLHa46
D
leus .
ng,Thyroid carcinoma,
tion:May play a role at early stages of development of specific neural lineages ost regions of the CNS, and of several lineages in the PNS. Essential for the eration of olfactory and autonomic neurons. Activates transcription by binding be E box (5'-CANNTG-3').,similarity:Contains 1 basic helix-loop-helix (bHLH) hain.,subunit:Efficient DNA binding requires dimerization with another bHLH ein. Forms a heterodimer with TCF3.,
aete-scute family bHLH transcription factor 1(ASCL1) Homo sapiens This e encodes a member of the basic helix-loop-helix (BHLH) family of scription factors. The protein activates transcription by binding to the E box apos;-CANNTG-3'). Dimerization with other BHLH proteins is required

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658




国内优质抗体供应商 精准的 WB 检测服务

24H 在线服务, 欢迎咨询

	for efficient DNA binding. This protein plays a role in the neuronal commitment and differentiation and in the generation of olfactory and autonomic neurons. Mutations in this gene may contribute to the congenital central hypoventilation syndrome (CCHS) phenotype in rare cases. [provided by RefSeq, Jul 2008],
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Immunohistochemistry analysis of ASCL1 antibody in paraffin-embedded human brain tissue.

	-117 -85
	-49
	-34
ASCL1-	- -25

Western blot analysis of lysate from Jurkat cells, using ASCL1 antibody.

Nanjing BYabscience technology Co.,Ltd