



HMG-17 Polyclonal Antibody

Catalog No	BYab-01794
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	IHC;IF;ELISA
Gene Name	HMGN2
Protein Name	Non-histone chromosomal protein HMG-17
Immunogen	The antiserum was produced against synthesized peptide derived from human HMG17. AA range:1-50
Specificity	HMG-17 Polyclonal Antibody detects endogenous levels of HMG-17 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/40000. Not yet tested in other applications.
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	HMGN2; HMG17; Non-histone chromosomal protein HMG-17; High mobility group nucleosome-binding domain-containing protein 2
Observed Band	15-17kD
Cell Pathway	Nucleus . Cytoplasm . Cytoplasmic enrichment upon phosphorylation.
Tissue Specificity	Brain,Colon,Eye,Lung,Lymph,Mammary gland,Muscle,Peripheral
Function	function: Binds to the inner side of the nucleosomal DNA thus altering the interaction between the DNA and the histone octamer. May be involved in the process which maintains transcribable genes in a unique chromatin conformation.,mass spectrometry: PubMed:10739259,PTM:Phosphorylation favors cytoplasmic localization.,similarity:Belongs to the HMGN family.,subcellular location:Cytoplasmic enrichment upon phosphorylation.,
Background	high mobility group nucleosomal binding domain 2(HMGN2) Homo sapiens The protein encoded by this gene binds nucleosomal DNA and is associated with transcriptionally active chromatin. Along with a similar protein, HMGN1, the

Nanjing BYabscience technology Co.,Ltd



encoded protein may help maintain an open chromatin configuration around transcribable genes. The protein has also been found to have antimicrobial activity against bacteria, viruses and fungi. [provided by RefSeq, Oct 2014],


matters needing attention

Avoid repeated freezing and thawing!


Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Immunofluorescence analysis of HeLa cells, using HMG17 Antibody. The picture on the right is blocked with the synthesized peptide.



Immunohistochemistry analysis of paraffin-embedded human brain tissue, using HMG17 Antibody. The picture on the right is blocked with the synthesized peptide.