



MafF Polyclonal Antibody

Catalog No	BYab-01856
Isotype	IgG
Reactivity	Human;Mouse
Applications	IHC;IF;ELISA
Gene Name	MAFF
Protein Name	Transcription factor MafF
Immunogen	The antiserum was produced against synthesized peptide derived from human MAFF. AA range:51-100
Specificity	MafF Polyclonal Antibody detects endogenous levels of MafF protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	IHC: 1/100 - 1/300. ELISA: 1/10000.. IF 1:50-200
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	MAFF; Transcription factor MafF; U-Maf; V-maf musculoaponeurotic fibrosarcoma oncogene homolog F
Observed Band	
Cell Pathway	Nucleus.
Tissue Specificity	Expressed in the term myometrium and kidney.
Function	function:Interacts with the upstream promoter region of the oxytocin receptor gene. May be a transcriptional enhancer in the up-regulation of the oxytocin receptor gene at parturition. Since it lacks a putative transactivation domain, it may behave as a transcriptional repressor when it dimerize among himself. May also serve as a transcriptional activator by dimerizing with other (usually larger) basic-zipper proteins and recruiting them to specific DNA-binding sites. May be involved in the cellular stress response.,induction:By oxidative stress.,similarity:Belongs to the bZIP family.,similarity:Belongs to the bZIP family. Maf subfamily.,similarity:Contains 1 bZIP domain.,subunit:Monomer and homo- or heterodimer. Interacts with MIP.,tissue specificity:Expressed in the term

Nanjing BYabscience technology Co.,Ltd



myometrium and kidney.,

Background

The protein encoded by this gene is a basic leucine zipper (bZIP) transcription factor that lacks a transactivation domain. It is known to bind the US-2 DNA element in the promoter of the oxytocin receptor (OTR) gene and most likely heterodimerizes with other leucine zipper-containing proteins to enhance expression of the OTR gene during term pregnancy. The encoded protein can also form homodimers, and since it lacks a transactivation domain, the homodimer may act as a repressor of transcription. This gene may also be involved in the cellular stress response. Multiple transcript variants encoding two different isoforms have been found for this gene. [provided by RefSeq, Jun 2009],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Immunohistochemistry analysis of paraffin-embedded human breast carcinoma tissue, using MAFF Antibody. The picture on the right is blocked with the synthesized peptide.