



# MN1 Polyclonal Antibody

<b>Catalog No</b>	BYab-01873
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Rat;Mouse;
<b>Applications</b>	WB;ELISA;IHC
<b>Gene Name</b>	MN1
<b>Protein Name</b>	Probable tumor suppressor protein MN1
<b>Immunogen</b>	The antiserum was produced against synthesized peptide derived from human MN1. AA range:821-870
<b>Specificity</b>	MN1 Polyclonal Antibody detects endogenous levels of MN1 protein.
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source</b>	Polyclonal, Rabbit,IgG
<b>Purification</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Dilution</b>	WB 1:500-2000;IHC-p 1:50-300; ELISA 2000-20000
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	MN1; Probable tumor suppressor protein MN1
<b>Observed Band</b>	135kD
<b>Cell Pathway</b>	Nucleus .
<b>Tissue Specificity</b>	Widely expressed in fetal and adult tissues. Highest expression is observed in fetal brain and skeletal muscle, and adult skeletal muscle.
<b>Function</b>	disease:A chromosomal aberration involving MN1 may be a cause of acute myeloid leukemia (AML). Translocation t(12;22)(p13;q11) with TEL.,disease:Defects in MN1 may be a cause of meningiomas, slowly growing benign tumors derived from the arachnoidal cap cells of the leptomeninges, the soft coverings of the brain and spinal cord. Meningiomas are believed to be the most common primary tumors of the central nervous system in man.,function:May play a role in tumor suppression.,tissue specificity:Ubiquitously expressed. Highest levels in skeletal muscle.,
<b>Background</b>	Meningioma 1 (MN1) contains two sets of CAG repeats. It is disrupted by a balanced translocation (4;22) in a meningioma, and its inactivation may contribute

Nanjing BYabscience technology Co.,Ltd



to meningioma 32 pathogenesis. [provided by RefSeq, Jul 2008],


**matters needing attention**

Avoid repeated freezing and thawing!


**Usage suggestions**

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

**Products Images**




Western Blot analysis of various cells using MN1 Polyclonal Antibody diluted at 1:2000



Western blot analysis of lysates from COLO cells, using MN1 Antibody. The lane on the right is blocked with the synthesized peptide.

Nanjing BYabscience technology Co.,Ltd



Immunohistochemical analysis of paraffin-embedded human liver cancer. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).