



NRIP2 Polyclonal Antibody

Catalog No BYab-01918 Isotype IgG Reactivity Human;Mouse Applications IHC;IF;ELISA Gene Name NRIP2 Protein Name Nuclear receptor-interacting protein 2 Immunogen The antiserum was produced against synthesized peptide derived from human NRIP2. AA range:1-50 Specificity NRIP2 Polyclonal Antibody detects endogenous levels of NRIP2 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution IHC: 1/100 - 1/300. ELISA: 1/20000 IF 1:50-200 Concentration 1 mg/ml Purity 290% Storage Stability -20°C/1 year Synonyms NRIP2; Nuclear receptor-interacting protein 2 Observed Band Cell Pathway Nucleus. Tissue Specificity Amygdala,Brain,PCR rescued clones,Trachea, MR1F2, subunit-interacts with NR1F2, RARA and THRB in a ligand-dependent manner. Background function:Down-regulates transcriptional activation by nuclear recep		
Reactivity Human;Mouse Applications IHC;IF;ELISA Gene Name NRIP2 Protein Name Nuclear receptor-interacting protein 2 Immunogen The antiserum was produced against synthesized peptide derived from human NRIP2. AA range:1-50 Specificity NRIP2 Polyclonal Antibody detects endogenous levels of NRIP2 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution IHC: 1/100 - 1/300. ELISA: 1/20000 IF 1:50-200 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms NRIP2; Nuclear receptor-interacting protein 2 Observed Band Cell Pathway Nucleus . Tissue Specificity Amygdala,Brain,PCR rescued clones,Trachea, function:Down-regulates transcriptional activation by nuclear receptors such as NR1F2. subunit:Interacts with NR1F2, RARA and THRB in a ligand-dependent manner., Background Varieties Avoid repeated freezing and thawing!	Catalog No	BYab-01918
Applications IHC;IF;ELISA Gene Name NRIP2 Protein Name Nuclear receptor-interacting protein 2 Immunogen The antiserum was produced against synthesized peptide derived from human NRIP2. AA range:1-50 Specificity NRIP2 Polyclonal Antibody detects endogenous levels of NRIP2 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution IHC: 1/100 - 1/300. ELISA: 1/20000 IF 1:50-200 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms NRIP2; Nuclear receptor-interacting protein 2 Observed Band Cell Pathway Nucleus . Tissue Specificity Amygdala,Brain,PCR rescued clones,Trachea, Function function:Down-regulates transcriptional activation by nuclear receptors such as NR1F2.,subunit:Interacts with NR1F2, RARA and THRB in a ligand-dependent manner., Background Avoid repeated freezing and thawing!	Isotype	IgG
Gene Name NRIP2 Protein Name Nuclear receptor-interacting protein 2 Immunogen The antiserum was produced against synthesized peptide derived from human NRIP2. AA range:1-50 Specificity NRIP2 Polyclonal Antibody detects endogenous levels of NRIP2 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution IHC: 1/100 - 1/300. ELISA: 1/20000 IF 1:50-200 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms NRIP2; Nuclear receptor-interacting protein 2 Observed Band Cell Pathway Nucleus . Tissue Specificity Amygdala,Brain,PCR rescued clones,Trachea, Function function:Down-regulates transcriptional activation by nuclear receptors such as NR1F2_subunit:Interacts with NR1F2, RARA and THRB in a ligand-dependent manner., Background Avoid repeated freezing and thawing!	Reactivity	Human;Mouse
Protein Name Nuclear receptor-interacting protein 2 Immunogen The antiserum was produced against synthesized peptide derived from human NRIP2. AA range:1-50 NRIP2 Polyclonal Antibody detects endogenous levels of NRIP2 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution IHC: 1/100 - 1/300. ELISA: 1/20000 IF 1:50-200 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms NRIP2; Nuclear receptor-interacting protein 2 Observed Band Cell Pathway Nucleus Tissue Specificity Amygdala, Brain, PCR rescued clones, Trachea, function: Down-regulates transcriptional activation by nuclear receptors such as NR1F2, subunit: Interacts with NR1F2, RARA and THRB in a ligand-dependent manner., Background function: Down-regulates transcriptional activation by nuclear receptors such as NR1F2, subunit: Interacts with NR1F2, RARA and THRB in a ligand-dependent manner., Matters needing Avoid repeated freezing and thawing!	Applications	IHC;IF;ELISA
Immunogen The antiserum was produced against synthesized peptide derived from human NRIP2. AA range:1-50 Specificity NRIP2 Polyclonal Antibody detects endogenous levels of NRIP2 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution IHC: 1/100 - 1/300. ELISA: 1/20000 IF 1:50-200 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms NRIP2; Nuclear receptor-interacting protein 2 Observed Band Cell Pathway Cell Pathway Nucleus . Tissue Specificity Amygdala,Brain,PCR rescued clones,Trachea, Function function:Down-regulates transcriptional activation by nuclear receptors such as NR1F2, subunit:Interacts with NR1F2, RARA and THRB in a ligand-dependent manner. Background function:Down-regulates transcriptional activation by nuclear receptors such as NR1F2, subunit:Interacts with NR1F2, RARA and THRB in a ligand-dependent manner. matters needing Avoid repeated freezing and thawing!	Gene Name	NRIP2
NRIP2. AA range:1-50 Specificity NRIP2 Polyclonal Antibody detects endogenous levels of NRIP2 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution IHC: 1/100 - 1/300. ELISA: 1/20000 IF 1:50-200 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms NRIP2; Nuclear receptor-interacting protein 2 Observed Band Cell Pathway Nucleus Tissue Specificity Amygdala,Brain,PCR rescued clones,Trachea, Function function:Down-regulates transcriptional activation by nuclear receptors such as NR1F2, subunit:Interacts with NR1F2, RARA and THRB in a ligand-dependent manner., Background function:Down-regulates transcriptional activation by nuclear receptors such as NR1F2, subunit:Interacts with NR1F2, RARA and THRB in a ligand-dependent manner., Matters needing Avoid repeated freezing and thawing!	Protein Name	Nuclear receptor-interacting protein 2
Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution IHC: 1/100 - 1/300. ELISA: 1/20000 IF 1:50-200 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms NRIP2; Nuclear receptor-interacting protein 2 Observed Band Cell Pathway Nucleus Tissue Specificity Amygdala, Brain, PCR rescued clones, Trachea, Function function: Down-regulates transcriptional activation by nuclear receptors such as NR1F2, subunit: Interacts with NR1F2, RARA and THRB in a ligand-dependent manner., Background function: Down-regulates transcriptional activation by nuclear receptors such as NR1F2, subunit: Interacts with NR1F2, RARA and THRB in a ligand-dependent manner., matters needing Avoid repeated freezing and thawing!	Immunogen	
Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution IHC: 1/100 - 1/300. ELISA: 1/20000 IF 1:50-200 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms NRIP2; Nuclear receptor-interacting protein 2 Observed Band Cell Pathway Nucleus . Tissue Specificity Amygdala, Brain, PCR rescued clones, Trachea, Function function: Down-regulates transcriptional activation by nuclear receptors such as NR1F2, subunit: Interacts with NR1F2, RARA and THRB in a ligand-dependent manner., Background function: Down-regulates transcriptional activation by nuclear receptors such as NR1F2, subunit: Interacts with NR1F2, RARA and THRB in a ligand-dependent manner., matters needing Avoid repeated freezing and thawing!	Specificity	NRIP2 Polyclonal Antibody detects endogenous levels of NRIP2 protein.
Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution IHC: 1/100 - 1/300. ELISA: 1/20000 IF 1:50-200 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms NRIP2; Nuclear receptor-interacting protein 2 Observed Band Cell Pathway Nucleus . Tissue Specificity Amygdala,Brain,PCR rescued clones,Trachea, Function function:Down-regulates transcriptional activation by nuclear receptors such as NR1F2.,subunit:Interacts with NR1F2, RARA and THRB in a ligand-dependent manner., Background function:Down-regulates transcriptional activation by nuclear receptors such as NR1F2.,subunit:Interacts with NR1F2, RARA and THRB in a ligand-dependent manner., Avoid repeated freezing and thawing!	Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen. Dilution IHC: 1/100 - 1/300. ELISA: 1/20000 IF 1:50-200 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms NRIP2; Nuclear receptor-interacting protein 2 Observed Band Cell Pathway Nucleus . Tissue Specificity Amygdala,Brain,PCR rescued clones,Trachea, Function function:Down-regulates transcriptional activation by nuclear receptors such as NR1F2.,subunit:Interacts with NR1F2, RARA and THRB in a ligand-dependent manner., Background function:Down-regulates transcriptional activation by nuclear receptors such as NR1F2.,subunit:Interacts with NR1F2, RARA and THRB in a ligand-dependent manner., Matters needing Avoid repeated freezing and thawing!	Source	Polyclonal, Rabbit,IgG
Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms NRIP2; Nuclear receptor-interacting protein 2 Observed Band Cell Pathway Nucleus . Tissue Specificity Amygdala,Brain,PCR rescued clones,Trachea, Function function:Down-regulates transcriptional activation by nuclear receptors such as NR1F2.,subunit:Interacts with NR1F2, RARA and THRB in a ligand-dependent manner., Background function:Down-regulates transcriptional activation by nuclear receptors such as NR1F2.,subunit:Interacts with NR1F2, RARA and THRB in a ligand-dependent manner., Matters needing Avoid repeated freezing and thawing!	Purification	
Purity ≥90% Storage Stability -20°C/1 year Synonyms NRIP2; Nuclear receptor-interacting protein 2 Observed Band Cell Pathway Nucleus . Tissue Specificity Amygdala,Brain,PCR rescued clones,Trachea, Function function:Down-regulates transcriptional activation by nuclear receptors such as NR1F2.,subunit:Interacts with NR1F2, RARA and THRB in a ligand-dependent manner., Background function:Down-regulates transcriptional activation by nuclear receptors such as NR1F2.,subunit:Interacts with NR1F2, RARA and THRB in a ligand-dependent manner., Matters needing Avoid repeated freezing and thawing!	Dilution	IHC: 1/100 - 1/300. ELISA: 1/20000 IF 1:50-200
Storage Stability -20°C/1 year Synonyms NRIP2; Nuclear receptor-interacting protein 2 Observed Band Cell Pathway Nucleus . Tissue Specificity Amygdala,Brain,PCR rescued clones,Trachea, Function function:Down-regulates transcriptional activation by nuclear receptors such as NR1F2,,subunit:Interacts with NR1F2, RARA and THRB in a ligand-dependent manner., Background function:Down-regulates transcriptional activation by nuclear receptors such as NR1F2,,subunit:Interacts with NR1F2, RARA and THRB in a ligand-dependent manner., matters needing Avoid repeated freezing and thawing!	Concentration	1 mg/ml
Synonyms NRIP2; Nuclear receptor-interacting protein 2 Observed Band Cell Pathway Nucleus . Tissue Specificity Amygdala,Brain,PCR rescued clones,Trachea, Function function:Down-regulates transcriptional activation by nuclear receptors such as NR1F2.,subunit:Interacts with NR1F2, RARA and THRB in a ligand-dependent manner., Background function:Down-regulates transcriptional activation by nuclear receptors such as NR1F2.,subunit:Interacts with NR1F2, RARA and THRB in a ligand-dependent manner., Matters needing Avoid repeated freezing and thawing!	Purity	≥90%
Observed Band Cell Pathway Nucleus . Tissue Specificity Amygdala,Brain,PCR rescued clones,Trachea, function:Down-regulates transcriptional activation by nuclear receptors such as NR1F2.,subunit:Interacts with NR1F2, RARA and THRB in a ligand-dependent manner., Background function:Down-regulates transcriptional activation by nuclear receptors such as NR1F2.,subunit:Interacts with NR1F2, RARA and THRB in a ligand-dependent manner., watters needing Avoid repeated freezing and thawing!	Storage Stability	-20°C/1 year
Cell Pathway Nucleus . Tissue Specificity Amygdala,Brain,PCR rescued clones,Trachea, function:Down-regulates transcriptional activation by nuclear receptors such as NR1F2.,subunit:Interacts with NR1F2, RARA and THRB in a ligand-dependent manner., Background function:Down-regulates transcriptional activation by nuclear receptors such as NR1F2.,subunit:Interacts with NR1F2, RARA and THRB in a ligand-dependent manner., matters needing Avoid repeated freezing and thawing!	Synonyms	NRIP2; Nuclear receptor-interacting protein 2
Tissue Specificity Amygdala,Brain,PCR rescued clones,Trachea, function:Down-regulates transcriptional activation by nuclear receptors such as NR1F2.,subunit:Interacts with NR1F2, RARA and THRB in a ligand-dependent manner., Background function:Down-regulates transcriptional activation by nuclear receptors such as NR1F2.,subunit:Interacts with NR1F2, RARA and THRB in a ligand-dependent manner., matters needing Avoid repeated freezing and thawing!	Observed Band	
Function function:Down-regulates transcriptional activation by nuclear receptors such as NR1F2.,subunit:Interacts with NR1F2, RARA and THRB in a ligand-dependent manner., function:Down-regulates transcriptional activation by nuclear receptors such as NR1F2.,subunit:Interacts with NR1F2, RARA and THRB in a ligand-dependent manner., matters needing Avoid repeated freezing and thawing!	Cell Pathway	Nucleus .
NR1F2.,subunit:Interacts with NR1F2, RARA and THRB in a ligand-dependent manner., Background function:Down-regulates transcriptional activation by nuclear receptors such as NR1F2.,subunit:Interacts with NR1F2, RARA and THRB in a ligand-dependent manner., matters needing Avoid repeated freezing and thawing!	Tissue Specificity	Amygdala,Brain,PCR rescued clones,Trachea,
NR1F2.,subunit:Interacts with NR1F2, RARA and THRB in a ligand-dependent manner., matters needing Avoid repeated freezing and thawing!	Function	NR1F2.,subunit:Interacts with NR1F2, RARA and THRB in a ligand-dependent
	Background	NR1F2.,subunit:Interacts with NR1F2, RARA and THRB in a ligand-dependent
	matters needing attention	Avoid repeated freezing and thawing!

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658




国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询



Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Immunohistochemistry analysis of paraffin-embedded human brain tissue, using NRIP2 Antibody. The picture on the right is blocked with the synthesized peptide.

Nanjing BYabscience technology Co.,Ltd