



# PRPF31 Polyclonal Antibody

<b>Catalog No</b>	BYab-01961
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Mouse
<b>Applications</b>	WB;ELISA;IHC
<b>Gene Name</b>	PRPF31
<b>Protein Name</b>	U4/U6 small nuclear ribonucleoprotein Prp31
<b>Immunogen</b>	The antiserum was produced against synthesized peptide derived from human PRPF31. AA range:331-380
<b>Specificity</b>	PRPF31 Polyclonal Antibody detects endogenous levels of PRPF31 protein.
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source</b>	Polyclonal, Rabbit,IgG
<b>Purification</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Dilution</b>	WB 1:500-2000;IHC-p 1:50-300; ELISA 2000-20000
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	PRPF31; PRP31; U4/U6 small nuclear ribonucleoprotein Prp31; Pre-mRNA-processing factor 31; Serologically defined breast cancer antigen NY-BR-99; U4/U6 snRNP 61 kDa protein; Protein 61K; hPrp31
<b>Observed Band</b>	55kD
<b>Cell Pathway</b>	Nucleus . Nucleus speckle . Nucleus, Cajal body . Predominantly found in speckles and in Cajal bodies. .
<b>Tissue Specificity</b>	Ubiquitously expressed.
<b>Function</b>	disease:Defects in PRPF31 are the cause of retinitis pigmentosa type 11 (RP11) [MIM:600138]. RP leads to degeneration of retinal photoreceptor cells. Patients typically have night vision blindness and loss of midperipheral visual field. As their condition progresses, they lose their far peripheral visual field and eventually central vision as well. RP11 inheritance is autosomal dominant.,function:Involved in pre-mRNA splicing. Required for U4/U6.U5 tri-snRNP formation.,similarity:Contains 1 Nop domain.,subcellular location:Predominantly found in speckles and in Cajal bodies.,subunit:Part of a tri-snRNP complex. Interacts with C20orf14/U5 snRNP-associated 102 kDa protein.,tissue

**Nanjing BYabscience technology Co.,Ltd**



specificity:Ubiquitously expressed.,

**Background**

This gene encodes a component of the spliceosome complex and is one of several retinitis pigmentosa-causing genes. When the gene product is added to the spliceosome complex, activation occurs.[provided by RefSeq, Jan 2009],


**matters needing attention**

Avoid repeated freezing and thawing!


**Usage suggestions**

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

**Products Images**



Western blot analysis of lysates from HeLa cells, using PRP31 Antibody. The lane on the right is blocked with the synthesized peptide.



Immunohistochemical analysis of paraffin-embedded human cervical carcinoma. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).