



## ZC3H13 Polyclonal Antibody

Catalog No         BYab-02165           Isotype         IgG           Reactivity         Human;Mouse           Applications         WB;ELISA           Gene Name         ZC3H13           Protein Name         Zinc finger CCCH domain-containing protein 13           Immunogen         The antiserum was produced against synthesized peptide derived from human ZC3H13. AA range:1571-1620           Specificity         ZC3H13 Polyclonal Antibody detects endogenous levels of ZC3H13 protein.           Formulation         Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.           Source         Polyclonal, Rabbit,IgG           Purification         The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.           Dilution         Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications.           Concentration         1 mg/ml           Purity         ≥90%           Storage Stability         -20°C/1 year           Synonyms         ZC3H13; KIAA0853; Zinc finger CCCH domain-containing protein 13           Observed Band         200kD           Cell Pathway         Nucleus speckle . Nucleus, nucleoplasm .           Tissue Specificity         Brain,Breast,Epithelium,Fetal brain,T-cell,Testis,Uterus,           Function         PTM:Phosphor		
Reactivity Human; Mouse  Applications WB; ELISA  Gene Name ZC3H13  Protein Name Zinc finger CCCH domain-containing protein 13  Immunogen The antiserum was produced against synthesized peptide derived from human ZC3H13. AA range:1571-1620  Specificity ZC3H13 Polyclonal Antibody detects endogenous levels of ZC3H13 protein.  Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  Source Polyclonal, Rabbit, IgG  Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications.  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms ZC3H13; KIAA0853; Zinc finger CCCH domain-containing protein 13  Observed Band 200kD  Cell Pathway Nucleus speckle . Nucleus, nucleoplasm .  Tissue Specificity Brain, Breast, Epithelium, Fetal brain, T-cell, Testis, Uterus,  Function PTM:Phosphorylated upon DNA damage, probably by ATM or ATR., similarity:Contains 1 C3H1-type zinc finger.,  Background Avoid repeated freezing and thawing!	Catalog No	BYab-02165
Applications WB;ELISA  Gene Name ZC3H13  Protein Name Zinc finger CCCH domain-containing protein 13  Immunogen The antiserum was produced against synthesized peptide derived from human ZC3H13. AA range:1571-1620  Specificity ZC3H13 Polyclonal Antibody detects endogenous levels of ZC3H13 protein.  Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  Source Polyclonal, Rabbit, IgG  Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications.  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms ZC3H13; KIAA0853; Zinc finger CCCH domain-containing protein 13  Observed Band 200kD  Cell Pathway Nucleus speckle . Nucleus, nucleoplasm .  Tissue Specificity Brain,Breast,Epithelium,Fetal brain,T-cell,Testis,Uterus,  Function PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity.Contains 1 C3H1-type zinc finger.,  Background Avoid repeated freezing and thawing!	Isotype	IgG
Gene Name         ZC3H13           Protein Name         Zinc finger CCCH domain-containing protein 13           Immunogen         The antiserum was produced against synthesized peptide derived from human ZC3H13. AA range:1571-1620           Specificity         ZC3H13 Polyclonal Antibody detects endogenous levels of ZC3H13 protein.           Formulation         Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.           Source         Polyclonal, Rabbit,lgG           Purification         The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.           Dilution         Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications.           Concentration         1 mg/ml           Purity         ≥90%           Storage Stability         -20°C/1 year           Synonyms         ZC3H13; KIAA0853; Zinc finger CCCH domain-containing protein 13           Observed Band         200kD           Cell Pathway         Nucleus speckle . Nucleus, nucleoplasm .           Tissue Specificity         Brain,Breast,Epithelium,Fetal brain,T-cell,Testis,Uterus,           Function         PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Contains 1 C3H1-type zinc finger.,           Background         Avoid repeated freezing and thawing!	Reactivity	Human;Mouse
Protein Name  Zinc finger CCCH domain-containing protein 13  Immunogen  The antiserum was produced against synthesized peptide derived from human ZC3H13. AA range:1571-1620  Specificity  ZC3H13 Polyclonal Antibody detects endogenous levels of ZC3H13 protein.  Formulation  Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  Source  Polyclonal, Rabbit,IgG  Purification  The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution  Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications.  Concentration  1 mg/ml  Purity  290%  Storage Stability  -20°C/1 year  Synonyms  ZC3H13; KIAA0853; Zinc finger CCCH domain-containing protein 13  Observed Band  200kD  Cell Pathway  Nucleus speckle . Nucleus, nucleoplasm .  Tissue Specificity  Brain,Breast,Epithelium,Fetal brain,T-cell,Testis,Uterus,  Function  PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Contains 1 C3H1-type zinc finger.,  Background  AVoid repeated freezing and thawing!	Applications	WB;ELISA
Immunogen       The antiserum was produced against synthesized peptide derived from human ZC3H13. AA range:1571-1620         Specificity       ZC3H13 Polyclonal Antibody detects endogenous levels of ZC3H13 protein.         Formulation       Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.         Source       Polyclonal, Rabbit,IgG         Purification       The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.         Dilution       Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications.         Concentration       1 mg/ml         Purity       ≥90%         Storage Stability       -20°C/1 year         Synonyms       ZC3H13; KIAA0853; Zinc finger CCCH domain-containing protein 13         Observed Band       200kD         Cell Pathway       Nucleus speckle . Nucleus, nucleoplasm .         Tissue Specificity       Brain,Breast,Epithelium,Fetal brain,T-cell,Testis,Uterus,         Function       PTM:Phosphorylated upon DNA damage, probably by ATM or ATR, similarity:Contains 1 C3H1-type zinc finger.,         Background       AYM:Phosphorylated upon DNA damage, probably by ATM or ATR, similarity:Contains 1 C3H1-type zinc finger.,         matters needing       Avoid repeated freezing and thawing!	Gene Name	ZC3H13
Specificity ZC3H13. AA range:1571-1620  Specificity ZC3H13 Polyclonal Antibody detects endogenous levels of ZC3H13 protein.  Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  Source Polyclonal, Rabbit,IgG  Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications.  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms ZC3H13; KIAA0853; Zinc finger CCCH domain-containing protein 13  Observed Band 200kD  Cell Pathway Nucleus speckle . Nucleus, nucleoplasm .  Tissue Specificity Brain,Breast,Epithelium,Fetal brain,T-cell,Testis,Uterus,  Function PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Contains 1 C3H1-type zinc finger.,  Background Avoid repeated freezing and thawing!	Protein Name	Zinc finger CCCH domain-containing protein 13
Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  Source Polyclonal, Rabbit, IgG  The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications.  Concentration 1 mg/ml  Purity ≥90% Storage Stability -20°C/1 year  Synonyms ZC3H13; KIAA0853; Zinc finger CCCH domain-containing protein 13  Observed Band 200kD  Cell Pathway Nucleus speckle . Nucleus, nucleoplasm .  Tissue Specificity Brain,Breast,Epithelium,Fetal brain,T-cell,Testis,Uterus,  Function PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Contains 1 C3H1-type zinc finger.,  Background Avoid repeated freezing and thawing!	lmmunogen	
Source       Polyclonal, Rabbit, IgG         Purification       The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.         Dilution       Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications.         Concentration       1 mg/ml         Purity       ≥90%         Storage Stability       -20°C/1 year         Synonyms       ZC3H13; KIAA0853; Zinc finger CCCH domain-containing protein 13         Observed Band       200kD         Cell Pathway       Nucleus speckle . Nucleus, nucleoplasm .         Tissue Specificity       Brain,Breast,Epithelium,Fetal brain,T-cell,Testis,Uterus,         Function       PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Contains 1 C3H1-type zinc finger.,         Background       PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Contains 1 C3H1-type zinc finger.,         matters needing       Avoid repeated freezing and thawing!	Specificity	ZC3H13 Polyclonal Antibody detects endogenous levels of ZC3H13 protein.
Purification  The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution  Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications.  Concentration  1 mg/ml  Purity  ≥90%  Storage Stability  -20°C/1 year  Synonyms  ZC3H13; KIAA0853; Zinc finger CCCH domain-containing protein 13  Observed Band  200kD  Cell Pathway  Nucleus speckle . Nucleus, nucleoplasm .  Tissue Specificity  Brain,Breast,Epithelium,Fetal brain,T-cell,Testis,Uterus,  Function  PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Contains 1 C3H1-type zinc finger.,  Background  PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Contains 1 C3H1-type zinc finger.,  matters needing  Avoid repeated freezing and thawing!	Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen.  Dilution Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications.  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms ZC3H13; KIAA0853; Zinc finger CCCH domain-containing protein 13  Observed Band 200kD  Cell Pathway Nucleus speckle . Nucleus, nucleoplasm .  Tissue Specificity Brain,Breast,Epithelium,Fetal brain,T-cell,Testis,Uterus,  Function PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Contains 1 C3H1-type zinc finger.,  Background PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Contains 1 C3H1-type zinc finger.,  matters needing Avoid repeated freezing and thawing!	Source	Polyclonal, Rabbit,IgG
applications.  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms ZC3H13; KIAA0853; Zinc finger CCCH domain-containing protein 13  Observed Band 200kD  Cell Pathway Nucleus speckle . Nucleus, nucleoplasm .  Tissue Specificity Brain,Breast,Epithelium,Fetal brain,T-cell,Testis,Uterus,  Function PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Contains 1 C3H1-type zinc finger.,  Background PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Contains 1 C3H1-type zinc finger.,  matters needing Avoid repeated freezing and thawing!	Purification	· · · · · · · · · · · · · · · · · · ·
Purity ≥90%  Storage Stability -20°C/1 year  Synonyms ZC3H13; KIAA0853; Zinc finger CCCH domain-containing protein 13  Observed Band 200kD  Cell Pathway Nucleus speckle . Nucleus, nucleoplasm .  Tissue Specificity Brain,Breast,Epithelium,Fetal brain,T-cell,Testis,Uterus,  Function PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Contains 1 C3H1-type zinc finger.,  Background PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Contains 1 C3H1-type zinc finger.,  matters needing Avoid repeated freezing and thawing!	Dilution	Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications.
Storage Stability  -20°C/1 year  ZC3H13; KIAA0853; Zinc finger CCCH domain-containing protein 13  Observed Band  200kD  Cell Pathway  Nucleus speckle . Nucleus, nucleoplasm .  Tissue Specificity  Brain,Breast,Epithelium,Fetal brain,T-cell,Testis,Uterus,  Function  PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Contains 1 C3H1-type zinc finger.,  Background  PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Contains 1 C3H1-type zinc finger.,  matters needing  Avoid repeated freezing and thawing!	Concentration	1 mg/ml
Synonyms  ZC3H13; KIAA0853; Zinc finger CCCH domain-containing protein 13  Observed Band  200kD  Cell Pathway  Nucleus speckle . Nucleus, nucleoplasm .  Tissue Specificity  Brain,Breast,Epithelium,Fetal brain,T-cell,Testis,Uterus,  Function  PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Contains 1 C3H1-type zinc finger.,  Background  PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Contains 1 C3H1-type zinc finger.,  matters needing  Avoid repeated freezing and thawing!	Purity	≥90%
Observed Band  Cell Pathway  Nucleus speckle . Nucleus, nucleoplasm .  Tissue Specificity  Brain,Breast,Epithelium,Fetal brain,T-cell,Testis,Uterus,  Function  PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Contains 1 C3H1-type zinc finger.,  Background  PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Contains 1 C3H1-type zinc finger.,  matters needing  Avoid repeated freezing and thawing!	Storage Stability	-20°C/1 year
Cell Pathway       Nucleus speckle . Nucleus, nucleoplasm .         Tissue Specificity       Brain,Breast,Epithelium,Fetal brain,T-cell,Testis,Uterus,         Function       PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Contains 1 C3H1-type zinc finger.,         Background       PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Contains 1 C3H1-type zinc finger.,         matters needing       Avoid repeated freezing and thawing!	Synonyms	ZC3H13; KIAA0853; Zinc finger CCCH domain-containing protein 13
Tissue Specificity  Brain,Breast,Epithelium,Fetal brain,T-cell,Testis,Uterus,  Function  PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Contains 1 C3H1-type zinc finger.,  PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Contains 1 C3H1-type zinc finger.,  matters needing  Avoid repeated freezing and thawing!	Observed Band	200kD
Function  PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Contains 1 C3H1-type zinc finger.,  PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Contains 1 C3H1-type zinc finger.,  matters needing  Avoid repeated freezing and thawing!	Cell Pathway	Nucleus speckle . Nucleus, nucleoplasm .
ATR.,similarity:Contains 1 C3H1-type zinc finger.,  Background PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Contains 1 C3H1-type zinc finger.,  matters needing Avoid repeated freezing and thawing!	Tissue Specificity	Brain,Breast,Epithelium,Fetal brain,T-cell,Testis,Uterus,
Background PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Contains 1 C3H1-type zinc finger.,  matters needing Avoid repeated freezing and thawing!	Function	PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Contains 1 C3H1-type zinc finger.,
	Background	PTM:Phosphorylated upon DNA damage, probably by ATM or
	matters needing attention	Avoid repeated freezing and thawing!

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658







**Usage suggestions** 

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.





Western Blot analysis of various cells using ZC3H13 Polyclonal Antibody. Secondary antibody(catalog#:RS0002) was diluted at 1:20000



Western blot analysis of lysates from A549, Jurkat, HepG2 cells, using ZC3H13 Antibody. The lane on the right is blocked with the synthesized peptide.

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658



国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询



