



ACOT8 Polyclonal Antibody

Catalog No	BYab-02472
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB;IHC;IF;ELISA
Gene Name	ACOT8
Protein Name	Acyl-coenzyme A thioesterase 8
Immunogen	The antiserum was produced against synthesized peptide derived from human ACOT8. AA range:131-180
Specificity	ACOT8 Polyclonal Antibody detects endogenous levels of ACOT8 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/40000. Not yet tested in other applications.
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	ACOT8; ACTEIII; PTE1; PTE2; Acyl-coenzyme A thioesterase 8; Acyl-CoA thioesterase 8; Choloyl-coenzyme A thioesterase; HIV-Nef-associated acyl-CoA thioesterase; PTE-2; Peroxisomal acyl-coenzyme A thioester hydrolase 1; PTE-1; Peroxisomal lon
Observed Band	36kD
Cell Pathway	Peroxisome matrix . Predominantly localized in the peroxisome but a localization to the cytosol cannot be excluded. .
Tissue Specificity	Detected in a T-cell line (at protein level). Ubiquitous (PubMed:9153233, PubMed:9299485).
Function	catalytic activity:Choloyl-CoA + H(2)O = cholate + CoA.,function:Acyl-CoA thioesterases are a group of enzymes that catalyze the hydrolysis of acyl-CoAs to the free fatty acid and coenzyme A (CoASH), providing the potential to regulate intracellular levels of acyl-CoAs, free fatty acids and CoASH. May mediate Nef-induced down-regulation of CD4. Major thioesterase in peroxisomes. Competes with BAAT (Bile acid CoA: amino acid N-acyltransferase) for bile

Nanjing BYabscience technology Co.,Ltd



acid-CoA substrate (such as chenodeoxycholoyl-CoA). Shows a preference for medium-length fatty acyl-CoAs (By similarity). May be involved in the metabolic regulation of peroxisome proliferation.,induction:Regulated by peroxisome proliferator (such as Clofibrate), via the peroxisome proliferator-activated receptors (PPARs),similarity:Belongs to the C/M/P thioester hydrolase family.,subunit:Interacts with HIV-1 Nef.,tissue specificity:Detect

Background

The protein encoded by this gene is a peroxisomal thioesterase that appears to be involved more in the oxidation of fatty acids rather than in their formation. The encoded protein can bind to the human immunodeficiency virus-1 protein Nef, and mediate Nef-induced down-regulation of CD4 in T-cells. [provided by RefSeq, Oct 2010],

matters needing attention


Avoid repeated freezing and thawing!

Usage suggestions


This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.




Products Images




Western Blot analysis of various cells using ACOT8 Polyclonal Antibody




Western Blot analysis of HuvEc cells using ACOT8 Polyclonal Antibody



Immunohistochemical analysis of paraffin-embedded Human brain. Antibody was diluted at 1:100(4° overnight). High-pressure and temperature Tris-EDTA,pH8.0 was used for antigen retrieval. Negative contrl (right) obtaned from antibody was pre-absorbed by immunogen peptide.



Immunofluorescence analysis of A549 cells, using ACOT8 Antibody. The picture on the right is blocked with the synthesized peptide.



Western blot analysis of lysates from HUVEC cells, using ACOT8 Antibody. The lane on the right is blocked with the synthesized peptide.