



DD2 Polyclonal Antibody

Catalog No	BYab-02611
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB;IHC;IF;ELISA
Gene Name	AKR1C2
Protein Name	Aldo-keto reductase family 1 member C2
Immunogen	The antiserum was produced against synthesized peptide derived from human AKR1C2. AA range:21-70
Specificity	DD2 Polyclonal Antibody detects endogenous levels of DD2 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB: 1/500 - 1/2000. IHC: 1/100 - 1/300. ELISA: 1/20000.. IF 1:50-200
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	AKR1C2; DDH2; Aldo-keto reductase family 1 member C2; 3-alpha-HSD3; Chlordecone reductase homolog HAKRD; Dihydrodiol dehydrogenase 2; DD-2; DD2; Dihydrodiol dehydrogenase/bile acid-binding protein; DD/BABP; Trans-1; 2-dihydrobenzene-1,2-diol
Observed Band	37kD
Cell Pathway	Cytoplasm, cytosol .
Tissue Specificity	Expressed in fetal testes. Expressed in fetal and adult adrenal glands.
Function	catalytic activity:Androsterone + NAD(P)(+) = 5-alpha-androstane-3,17-dione + NAD(P)H.,catalytic activity:Trans-1,2-dihydrobenzene-1,2-diol + NADP(+) = catechol + NADPH.,enzyme regulation:Inhibited by hexestrol with an IC(50) of 2.8 uM, 1,10-phenanthroline with an IC(50) of 2100 uM, 1,7-phenanthroline with an IC(50) of 1500 uM, flufenamic acid with an IC(50) of 0.9 uM, indomethacin with an IC(50) of 75 uM, ibuprofen with an IC(50) of 6.9 uM, lithocholic acid with an IC(50) of 0.07 uM, ursodeoxycholic acid with an IC(50) of 0.08 uM and chenodeoxycholic

Nanjing BYabscience technology Co.,Ltd



acid with an IC(50) of 0.13 uM.,function:Works in concert with the 5-alpha/5-beta-steroid reductases to convert steroid hormones into the 3-alpha/5-alpha and 3-alpha/5-beta-tetrahydrosteroids. Catalyzes the inactivation of the most potent androgen 5-alpha-dihydrotestosterone (5-alpha-DHT) to 5-alpha-androstane-3-alpha,17-beta-diol (3-alpha-

Background

This gene encodes a member of the aldo/keto reductase superfamily, which consists of more than 40 known enzymes and proteins. These enzymes catalyze the conversion of aldehydes and ketones to their corresponding alcohols using NADH and/or NADPH as cofactors. The enzymes display overlapping but distinct substrate specificity. This enzyme binds bile acid with high affinity, and shows minimal 3-alpha-hydroxysteroid dehydrogenase activity. This gene shares high sequence identity with three other gene members and is clustered with those three genes at chromosome 10p15-p14. Three transcript variants encoding two different isoforms have been found for this gene. [provided by RefSeq, Dec 2011],

matters needing attention

Avoid repeated freezing and thawing!


Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.


Nanjing BYabscience technology Co.,Ltd




Products Images



Western Blot analysis of COLO cells using DD2 Polyclonal Antibody



Western blot analysis of lysates from COLO cells, using AKR1C2 Antibody. The lane on the right is blocked with the synthesized peptide.



Immunohistochemical analysis of paraffin-embedded human lung cancer. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).