



FX Polyclonal Antibody

Catalog No	BYab-02637
Isotype	lgG
Reactivity	Human;Mouse;Rat
Applications	WB;ELISA
Gene Name	TSTA3
Protein Name	GDP-L-fucose synthase
Immunogen	The antiserum was produced against synthesized peptide derived from human TSTA3. AA range:221-270
Specificity	FX Polyclonal Antibody detects endogenous levels of FX protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	Western Blot: 1/500 - 1/2000. ELISA: 1/20000. Not yet tested in other applications.
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	TSTA3; SDR4E1; GDP-L-fucose synthase; GDP-4-keto-6-deoxy-D-mannose-3; 5-epimerase-4-reductase; Protein FX; Red cell NADP(H)-binding protein; Short-chain dehydrogenase/reductase family 4E member 1
Observed Band	40kD
Cell Pathway	cytoplasm,cytosol,extracellular exosome,
Tissue Specificity	Erythrocyte,Lung,Placenta,Skin,
Function	catalytic activity:GDP-L-fucose + NADP(+) = GDP-4-dehydro-6-deoxy-D-mannose + NADPH.,function:Two step NADP-dependent conversion of GDP-4-dehydro-6-deoxy-D-mannose to GDP-fucose, involving an epimerase and a reductase reaction.,pathway:Nucleotide-sugar biosynthesis; GDP-L-fucose biosynthesis via de novo pathway; GDP-L-fucose from GDP-D-mannose: step 2/2.,similarity:Belongs to the fucose synthetase family.,subunit:Homodimer.,

Nanjing BYabscience technology Co.,Ltd



48-

34-

26-

19-

国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务, 欢迎咨询



Tissue specific transplantation antigen P35B is a NADP(H)-binding protein. It catalyze the two-step epimerase and the reductase reactions in GDP-D-mannose metabolism, converting GDP-4-keto-6-D-deoxymannose to GDP-L-fucose. GDP-L-fucose is the substrate of several fucosyltransferases involved in the expression of many glycoconjugates, including blood group ABH antigens and developmental adhesion antigens. Mutations in this gene may cause leukocyte adhesion deficiency, type II. [provided by RefSeq, Jul 2008],
Avoid repeated freezing and thawing!
This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images muscle kidney muscle Western blot analysis of lysates from rat kidney and rat -- 117 muscle cells, using TSTA3 Antibody. The lane on the right is blocked with the synthesized peptide. -- 85 -- 48 TSTA3---- 34 -- 26 -- 19 (kD) (kD) Western blot analysis of the lysates from COLO205 cells using TSTA3 antibody. 117-85-TSTA3

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658