



# MAN1B1 Polyclonal Antibody

<b>Catalog No</b>	BYab-02666
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Rat;Mouse;
<b>Applications</b>	WB;ELISA
<b>Gene Name</b>	MAN1B1
<b>Protein Name</b>	Endoplasmic reticulum mannosyl-oligosaccharide 1,2-alpha-mannosidase
<b>Immunogen</b>	Synthesized peptide derived from MAN1B1 . at AA range: 100-180
<b>Specificity</b>	MAN1B1 Polyclonal Antibody detects endogenous levels of MAN1B1 protein.
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source</b>	Polyclonal, Rabbit,IgG
<b>Purification</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Dilution</b>	Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications.
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	MAN1B1; Endoplasmic reticulum mannosyl-oligosaccharide 1; 2-alpha-mannosidase; ER alpha-1,2-mannosidase; ER mannosidase 1; ERMan1; Man9GlcNAc2-specific-processing alpha-mannosidase; Mannosidase alpha class 1B member 1
<b>Observed Band</b>	80kD
<b>Cell Pathway</b>	Endoplasmic reticulum membrane ; Single-pass type II membrane protein .
<b>Tissue Specificity</b>	Widely expressed.
<b>Function</b>	catalytic activity:Hydrolysis of the terminal (1->2)-linked alpha-D-mannose residues in the oligo-mannose oligosaccharide Man(9)(GlcNAc)(2).,cofactor:Calcium.,enzyme regulation:Inhibited by both 1-deoxymannojirimycin and kifunensine.,function:Involved in the maturation of Asn-linked oligosaccharides. Trim a single alpha-1,2-linked mannose residue from Man(9)GlcNAc(2) to produce Man(8)GlcNAc(2). The only product is the Man(8)GlcNAc(2) isomer B, the form lacking the middle-arm terminal alpha 1,2-mannose. It may be involved in glycoprotein quality control since it is

Nanjing BYabscience technology Co.,Ltd



important to target misfolded glycoproteins for degradation.,pathway:Protein modification; protein glycosylation.,similarity:Belongs to the glycosyl hydrolase 47 family.,tissue specificity:Widely expressed.,

**Background**

This gene encodes an enzyme belonging to the glycosyl hydrolase 47 family. This enzyme functions in N-glycan biosynthesis, and is a class I alpha-1,2-mannosidase that specifically converts Man9GlcNAc to Man8GlcNAc isomer B. It is required for N-glycan trimming to Man5-6GlcNAc2 in the endoplasmic-reticulum-associated degradation pathway. Mutations in this gene cause autosomal-recessive intellectual disability. Alternative splicing results in multiple transcript variants. A related pseudogene has been identified on chromosome 11. [provided by RefSeq, Dec 2011],

**matters needing attention**


Avoid repeated freezing and thawing!

**Usage suggestions**


This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.



## Products Images



Western Blot analysis of 3T3 cells using MAN1B1 Polyclonal Antibody



Western blot analysis of MAN1B1 Antibody. The lane on the right is blocked with the MAN1B1 peptide.