



PSMD12 Polyclonal Antibody

Catalog No	BYab-02768
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB;IHC;IF;ELISA
Gene Name	PSMD12
Protein Name	26S proteasome non-ATPase regulatory subunit 12
Immunogen	The antiserum was produced against synthesized peptide derived from human PSMD12. AA range:151-200
Specificity	PSMD12 Polyclonal Antibody detects endogenous levels of PSMD12 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB: 1/500 - 1/2000. IHC: 1/100 - 1/300. ELISA: 1/40000.. IF 1:50-200
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	PSMD12; 26S proteasome non-ATPase regulatory subunit 12; 26S proteasome regulatory subunit RPN5; 26S proteasome regulatory subunit p55
Observed Band	50kD
Cell Pathway	proteasome complex,nucleoplasm,cytoplasm,cytosol,proteasome regulatory particle,proteasome regulatory particle, lid subcomplex,membrane,proteasome accessory complex,nuclear proteasome complex,extracellular exosome,
Tissue Specificity	Brain,Muscle,Platelet,Testis,Tongue,
Function	function:Acts as a regulatory subunit of the 26S proteasome which is involved in the ATP-dependent degradation of ubiquitinated proteins.,similarity:Belongs to the proteasome subunit p55 family.,similarity:Contains 1 PCI domain.,subunit:Component of the PA700 complex.,
Background	The 26S proteasome is a multicatalytic proteinase complex with a highly ordered structure composed of 2 complexes, a 20S core and a 19S regulator. The 20S core is composed of 4 rings of 28 non-identical subunits; 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. The 19S regulator

Nanjing BYabscience technology Co.,Ltd



is composed of a base, which contains 6 ATPase subunits and 2 non-ATPase subunits, and a lid, which contains up to 10 non-ATPase subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. An essential function of a modified proteasome, the immunoproteasome, is the processing of class I MHC peptides. This gene encodes a non-ATPase subunit of the 19S regulator. A pseudogene has been identified on chromosome 3. Alternatively spliced transcript variants encoding


matters needing attention

Avoid repeated freezing and thawing!


Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images




Western Blot analysis of various cells using PSMD12 Polyclonal Antibody




Western Blot analysis of HepG2 cells using PSMD12 Polyclonal Antibody


Nanjing BYabscience technology Co.,Ltd



Western blot analysis of lysates from HepG2 cells, using PSD12 Antibody. The lane on the right is blocked with the synthesized peptide.



Western blot analysis of the lysates from HT-29 cells using PSMD12 antibody.



Immunohistochemical analysis of paraffin-embedded human tonsil. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 30min).