



USP42 Polyclonal Antibody

Catalog No BYab-02837 Isotype IgG Reactivity Human;Mouse;Rat Applications WB;ELISA Gene Name USP42 Protein Name Ubiquitin carboxyl-terminal hydrolase 42 Immunogen The antiserum was produced against synthesized peptide derived from human USP42. AA range:251-300 Specificity USP42 Polyclonal Antibody detects endogenous levels of USP42 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications. Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms USP42; Ubiquitin carboxyl-terminal hydrolase 42; Deubiquitinating enzyme 42; Ubiquitin thioesterase 42; Ubiquitin-specific-processing protease 42 Observed Band 150kD Cell Pathway Tissue Specificity Broadly expressed. Function catalytic activity: Ubiquitin C-ter		
Reactivity Human;Mouse;Rat Applications WB;ELISA Gene Name USP42 Protein Name Ubiquitin carboxyl-terminal hydrolase 42 Immunogen The antiserum was produced against synthesized peptide derived from human USP42. AA range:251-300 Specificity USP42 Polyclonal Antibody detects endogenous levels of USP42 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications. Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms USP42; Ubiquitin carboxyl-terminal hydrolase 42; Deubiquitinating enzyme 42; Ubiquitin thioesterase 42; Ubiquitin-specific-processing protease 42 Observed Band 150kD Cell Pathway Tissue Specificity Broadly expressed. Function catalytic activity: Ubiquitin C-terminal thioester + H(2)O = ubiquitin + a thiol, similarity: Belongs to the peptidase C19 family., tissue specificity: Broadly expressed. Background Avoid repeated freezing and thawing!	Catalog No	BYab-02837
Applications WB;ELISA Gene Name USP42 Protein Name Ubiquitin carboxyl-terminal hydrolase 42 Immunogen The antiserum was produced against synthesized peptide derived from human USP42. AA range:251-300 Specificity USP42 Polyclonal Antibody detects endogenous levels of USP42 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit.lgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications. Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms USP42; Ubiquitin carboxyl-terminal hydrolase 42; Deubiquitinating enzyme 42; Ubiquitin thioesterase 42; Ubiquitin-specific-processing protease 42 Observed Band 150kD Cell Pathway Tissue Specificity Broadly expressed. Function catalytic activity-Ubiquitin C-terminal thioester + H(2)O = ubiquitin + a thiol, similanty-Belongs to the peptidase C19 familly. tissue specificity:Broadly expressed. Background Avoid repeated freezing and thawing!	Isotype	IgG
Gene Name USP42 Protein Name Ubiquitin carboxyl-terminal hydrolase 42 Immunogen The antiserum was produced against synthesized peptide derived from human USP42. AA range:251-300 Specificity USP42 Polyclonal Antibody detects endogenous levels of USP42 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications. Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms USP42; Ubiquitin carboxyl-terminal hydrolase 42; Deubiquitinating enzyme 42; Ubiquitin thioesterase 42; Ubiquitin-specific-processing protease 42 Observed Band 150kD Cell Pathway Tissue Specificity Tissue Specificity Broadly expressed. Function catalytic activity: Ubiquitin C-terminal thioester + H(2)O = ubiquitin + a thiol., similarity. Belongs to the peptidase C19 family., tissue specificity:Broadly expressed. Background Avoid repeated freezing and thawing! <th>Reactivity</th> <th>Human;Mouse;Rat</th>	Reactivity	Human;Mouse;Rat
Protein Name Ubiquitin carboxyl-terminal hydrolase 42 Immunogen The antiserum was produced against synthesized peptide derived from human USP42. AA range:251-300 Specificity USP42 Polyclonal Antibody detects endogenous levels of USP42 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications. Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms USP42; Ubiquitin carboxyl-terminal hydrolase 42; Deubiquitinating enzyme 42; Ubiquitin thioesterase 42; Ubiquitin-specific-processing protease 42 Observed Band 150kD Cell Pathway Tissue Specificity Broadly expressed. Function catalytic activity: Ubiquitin C-terminal thioester + H(2)O = ubiquitin + a thiol., similarity: Belongs to the peptidase C19 family., tissue specificity: Broadly expressed. Background catalytic activity: Ubiquitin C-terminal thioester + H(2)O = ubiquitin + a thiol., similarity: Belongs to the peptidase C19 family., tissue specificity: Broadly expressed. Matters needing Avoid repeated freezing and thawing!	Applications	WB;ELISA
Immunogen The antiserum was produced against synthesized peptide derived from human USP42. AA range:251-300 Specificity USP42 Polyclonal Antibody detects endogenous levels of USP42 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications. Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms USP42; Ubiquitin carboxyl-terminal hydrolase 42; Deubiquitinating enzyme 42; Ubiquitin thioesterase 42; Ubiquitin-specific-processing protease 42 Observed Band 150kD Cell Pathway Tissue Specificity Broadly expressed. Function catalytic activity: Ubiquitin C-terminal thioester + H(2)O = ubiquitin + a thiol., similarity: Belongs to the peptidase C19 family., tissue specificity: Broadly expressed. Background catalytic activity: Ubiquitin C-terminal thioester + H(2)O = ubiquitin + a thiol., similarity: Belongs to the peptidase C19 family., tissue specificity: Broadly expressed. Avoid repeated freezing and thawing!	Gene Name	USP42
USP42. AA range:251-300 Specificity USP42 Polyclonal Antibody detects endogenous levels of USP42 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications. Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms USP42; Ubiquitin carboxyl-terminal hydrolase 42; Deubiquitinating enzyme 42; Ubiquitin thioesterase 42; Ubiquitin-specific-processing protease 42 Observed Band 150kD Cell Pathway Tissue Specificity Broadly expressed. Function catalytic activity: Ubiquitin C-terminal thioester + H(2)O = ubiquitin + a thiol., similarity: Belongs to the peptidase C19 family., tissue specificity: Broadly expressed., Background Avoid repeated freezing and thawing!	Protein Name	Ubiquitin carboxyl-terminal hydrolase 42
Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications. Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms USP42; Ubiquitin carboxyl-terminal hydrolase 42; Deubiquitinating enzyme 42; Ubiquitin thioesterase 42; Ubiquitin-specific-processing protease 42 Observed Band 150kD Cell Pathway Tissue Specificity Broadly expressed. Function catalytic activity: Ubiquitin C-terminal thioester + H(2)O = ubiquitin + a thiol., similarity: Belongs to the peptidase C19 family., tissue specificity: Broadly expressed., matters needing Avoid repeated freezing and thawing!	Immunogen	
Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications. Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms USP42; Ubiquitin carboxyl-terminal hydrolase 42; Deubiquitinating enzyme 42; Ubiquitin thioesterase 42; Ubiquitin-specific-processing protease 42 Observed Band 150kD Cell Pathway Tissue Specificity Broadly expressed. Function catalytic activity: Ubiquitin C-terminal thioester + H(2)O = ubiquitin + a thiol., similarity: Belongs to the peptidase C19 family., tissue specificity: Broadly expressed. Background catalytic activity: Ubiquitin C-terminal thioester + H(2)O = ubiquitin + a thiol., similarity: Belongs to the peptidase C19 family., tissue specificity: Broadly expressed. Matters needing Avoid repeated freezing and thawing!	Specificity	USP42 Polyclonal Antibody detects endogenous levels of USP42 protein.
Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications. Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms USP42; Ubiquitin carboxyl-terminal hydrolase 42; Deubiquitinating enzyme 42; Ubiquitin thioesterase 42; Ubiquitin-specific-processing protease 42 Observed Band 150kD Cell Pathway Tissue Specificity Broadly expressed. Function catalytic activity: Ubiquitin C-terminal thioester + H(2)O = ubiquitin + a thiol., similarity: Belongs to the peptidase C19 family., tissue specificity: Broadly expressed. Background catalytic activity: Ubiquitin C-terminal thioester + H(2)O = ubiquitin + a thiol., similarity: Belongs to the peptidase C19 family., tissue specificity: Broadly expressed. matters needing Avoid repeated freezing and thawing!	Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen. Dilution Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications. Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms USP42; Ubiquitin carboxyl-terminal hydrolase 42; Deubiquitinating enzyme 42; Ubiquitin thioesterase 42; Ubiquitin-specific-processing protease 42 Observed Band 150kD Cell Pathway Tissue Specificity Broadly expressed. Function catalytic activity: Ubiquitin C-terminal thioester + H(2)O = ubiquitin + a thiol., similarity: Belongs to the peptidase C19 family., tissue specificity: Broadly expressed., Background catalytic activity: Ubiquitin C-terminal thioester + H(2)O = ubiquitin + a thiol., similarity: Belongs to the peptidase C19 family., tissue specificity: Broadly expressed., Matters needing Avoid repeated freezing and thawing!	Source	Polyclonal, Rabbit,IgG
applications. Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms USP42; Ubiquitin carboxyl-terminal hydrolase 42; Deubiquitinating enzyme 42; Ubiquitin thioesterase 42; Ubiquitin-specific-processing protease 42 Observed Band 150kD Cell Pathway Tissue Specificity Broadly expressed. Function catalytic activity:Ubiquitin C-terminal thioester + H(2)O = ubiquitin + a thiol.,similarity:Belongs to the peptidase C19 family.,tissue specificity:Broadly expressed., Background catalytic activity:Ubiquitin C-terminal thioester + H(2)O = ubiquitin + a thiol.,similarity:Belongs to the peptidase C19 family.,tissue specificity:Broadly expressed., Matters needing Avoid repeated freezing and thawing!	Purification	·
Purity ≥90% Storage Stability -20°C/1 year Synonyms USP42; Ubiquitin carboxyl-terminal hydrolase 42; Deubiquitinating enzyme 42; Ubiquitin thioesterase 42; Ubiquitin-specific-processing protease 42 Observed Band 150kD Cell Pathway Tissue Specificity Broadly expressed. Function catalytic activity:Ubiquitin C-terminal thioester + H(2)O = ubiquitin + a thiol.,similarity:Belongs to the peptidase C19 family.,tissue specificity:Broadly expressed., Background catalytic activity:Ubiquitin C-terminal thioester + H(2)O = ubiquitin + a thiol.,similarity:Belongs to the peptidase C19 family.,tissue specificity:Broadly expressed., matters needing Avoid repeated freezing and thawing!	Dilution	Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications.
Storage Stability -20°C/1 year Synonyms USP42; Ubiquitin carboxyl-terminal hydrolase 42; Deubiquitinating enzyme 42; Ubiquitin thioesterase 42; Ubiquitin-specific-processing protease 42 Observed Band 150kD Cell Pathway Tissue Specificity Broadly expressed. Function catalytic activity: Ubiquitin C-terminal thioester + H(2)O = ubiquitin + a thiol., similarity: Belongs to the peptidase C19 family., tissue specificity: Broadly expressed., Background catalytic activity: Ubiquitin C-terminal thioester + H(2)O = ubiquitin + a thiol., similarity: Belongs to the peptidase C19 family., tissue specificity: Broadly expressed., Matters needing Avoid repeated freezing and thawing!	Concentration	1 mg/ml
Synonyms USP42; Ubiquitin carboxyl-terminal hydrolase 42; Deubiquitinating enzyme 42; Ubiquitin thioesterase 42; Ubiquitin-specific-processing protease 42 Observed Band 150kD Cell Pathway Tissue Specificity Broadly expressed. Function catalytic activity: Ubiquitin C-terminal thioester + H(2)O = ubiquitin + a thiol., similarity: Belongs to the peptidase C19 family., tissue specificity: Broadly expressed., Background catalytic activity: Ubiquitin C-terminal thioester + H(2)O = ubiquitin + a thiol., similarity: Belongs to the peptidase C19 family., tissue specificity: Broadly expressed., matters needing Avoid repeated freezing and thawing!	Purity	≥90%
Ubiquitin thioesterase 42; Ubiquitin-specific-processing protease 42 Observed Band 150kD Cell Pathway Tissue Specificity Broadly expressed. Function catalytic activity: Ubiquitin C-terminal thioester + H(2)O = ubiquitin + a thiol., similarity: Belongs to the peptidase C19 family., tissue specificity: Broadly expressed., Background catalytic activity: Ubiquitin C-terminal thioester + H(2)O = ubiquitin + a thiol., similarity: Belongs to the peptidase C19 family., tissue specificity: Broadly expressed., matters needing Avoid repeated freezing and thawing!	Storage Stability	-20°C/1 year
Cell Pathway Tissue Specificity Broadly expressed. Function catalytic activity:Ubiquitin C-terminal thioester + H(2)O = ubiquitin + a thiol.,similarity:Belongs to the peptidase C19 family.,tissue specificity:Broadly expressed., Background catalytic activity:Ubiquitin C-terminal thioester + H(2)O = ubiquitin + a thiol.,similarity:Belongs to the peptidase C19 family.,tissue specificity:Broadly expressed., matters needing Avoid repeated freezing and thawing!	Synonyms	
Tissue Specificity Broadly expressed. Function catalytic activity:Ubiquitin C-terminal thioester + H(2)O = ubiquitin + a thiol.,similarity:Belongs to the peptidase C19 family.,tissue specificity:Broadly expressed., Background catalytic activity:Ubiquitin C-terminal thioester + H(2)O = ubiquitin + a thiol.,similarity:Belongs to the peptidase C19 family.,tissue specificity:Broadly expressed., matters needing Avoid repeated freezing and thawing!	Observed Band	150kD
Function catalytic activity:Ubiquitin C-terminal thioester + H(2)O = ubiquitin + a thiol.,similarity:Belongs to the peptidase C19 family.,tissue specificity:Broadly expressed., Background catalytic activity:Ubiquitin C-terminal thioester + H(2)O = ubiquitin + a thiol.,similarity:Belongs to the peptidase C19 family.,tissue specificity:Broadly expressed., matters needing Avoid repeated freezing and thawing!	Cell Pathway	
thiol., similarity: Belongs to the peptidase C19 family., tissue specificity: Broadly expressed., Background catalytic activity: Ubiquitin C-terminal thioester + H(2)O = ubiquitin + a thiol., similarity: Belongs to the peptidase C19 family., tissue specificity: Broadly expressed., matters needing Avoid repeated freezing and thawing!	Tissue Specificity	Broadly expressed.
thiol., similarity: Belongs to the peptidase C19 family., tissue specificity: Broadly expressed., matters needing Avoid repeated freezing and thawing!	Function	thiol. similarity:Belongs to the peptidase C19 family, tissue specificity:Broadly
	Background	thiol., similarity: Belongs to the peptidase C19 family., tissue specificity: Broadly
	matters needing attention	Avoid repeated freezing and thawing!

Nanjing BYabscience technology Co.,Ltd


网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658





Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.





国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询



