




## TCP-1 α Polyclonal Antibody

Catalog No	BYab-03192	
Isotype	IgG	
Reactivity	Human;Mouse;Rat	
Applications	WB;ELISA	
Gene Name	TCP1	
Protein Name	T-complex protein 1 subunit alpha	
Immunogen	Synthesized peptide derived from the Internal region of human TCP-1 $\alpha$ .	
Specificity	TCP-1 $\alpha$ Polyclonal Antibody detects endogenous levels of TCP-1 $\alpha$ protein.	
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.	
Source	Polyclonal, Rabbit,IgG	
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.	
Dilution	Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications.	
Concentration	1 mg/ml	
Purity	≥90%	
Storage Stability	-20°C/1 year	
Synonyms	TCP1; CCT1; CCTA; T-complex protein 1 subunit alpha; TCP-1-alpha; CCT-alpha	
Observed Band	60kD	
Cell Pathway	Cytoplasm, cytosol . Cytoplasm, cytoskeleton, microtubule organizing center, centrosome .	
Tissue Specificity	Brain, Cajal-Retzius cell, Colon, Epithelium, Platelet, Testis,	
Function	function:Molecular chaperone; assists the folding of proteins upon ATP hydrolysis. Known to play a role, in vitro, in the folding of actin and tubulin.,similarity:Belongs to the TCP-1 chaperonin family.,subunit:Heterooligomeric complex of about 850 to 900 kDa that forms two stacked rings, 12 to 16 nm in diameter. Interacts with PACRG.,	
Background	The protein encoded by this gene is a molecular chaperone that is a member of the chaperonin containing TCP1 complex (CCT), also known as the TCP1 ring complex (TRiC). This complex consists of two identical stacked rings, each containing eight different proteins. Unfolded polypeptides enter the central cavity of the complex and are folded in an ATP-dependent manner. The complex folds various proteins, including actin and tubulin. Alternate transcriptional splice	

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 监督电话: 15950492658 官方热线: 025-5229-8998







	variants of this gene, encoding different isoforms, have been characterized. In addition, three pseudogenes that appear to be derived from this gene have been found. [provided by RefSeq, Jun 2010],	
matters needing attention	Avoid repeated freezing and thawing!	
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.	

Products Images				

Nanjing BYabscience technology Co.,Ltd