



BCAR3 Polyclonal Antibody

Catalog No	BYab-03740
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB;IHC;IF;ELISA
Gene Name	BCAR3
Protein Name	Breast cancer anti-estrogen resistance protein 3
Immunogen	The antiserum was produced against synthesized peptide derived from human BCAR3. AA range:761-810
Specificity	BCAR3 Polyclonal Antibody detects endogenous levels of BCAR3 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/10000. Not yet tested in other applications.
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	BCAR3; NSP2; SH2D3B; Breast cancer anti-estrogen resistance protein 3; Novel SH2-containing protein 2; SH2 domain-containing protein 3B
Observed Band	92kD
Cell Pathway	Cytoplasm . Cell junction, focal adhesion . Localization to focal adhesions depends on interaction with PTPRA. .
Tissue Specificity	Ubiquitously expressed. Found in several cancer cell lines, but not in nonmalignant breast tissue.
Function	function:May act as an adapter protein and couple activated growth factor receptors to a signaling pathway that regulates the proliferation in breast cancer cells. When overexpressed, it confers anti-estrogen resistance in breast cancer cell lines. May also be regulated by cellular adhesion to extracellular matrix proteins.,PTM:Phosphorylated on tyrosine.,similarity:Contains 1 Ras-GEF domain.,similarity:Contains 1 SH2 domain.,subunit:Interacts with BCAR1, NEDD9, PTK2 and PTPN1.,tissue specificity:Ubiquitously expressed. Found in several cancer cell lines, but not in nonmalignant breast tissue.,

Nanjing BYabscience technology Co.,Ltd



Background

breast cancer anti-estrogen resistance 3(BCAR3) Homo sapiens Breast tumors are initially dependent on estrogens for growth and progression and can be inhibited by anti-estrogens such as tamoxifen. However, breast cancers progress to become anti-estrogen resistant. Breast cancer anti-estrogen resistance gene 3 was identified in the search for genes involved in the development of estrogen resistance. The gene encodes a component of intracellular signal transduction that causes estrogen-independent proliferation in human breast cancer cells. The protein contains a putative src homology 2 (SH2) domain, a hall mark of cellular tyrosine kinase signaling molecules, and is partly homologous to the cell division cycle protein CDC48. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, May 2012],

matters needing attention


Avoid repeated freezing and thawing!

Usage suggestions


This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.




Products Images



Immunohistochemical analysis of paraffin-embedded Human breast cancer. Antibody was diluted at 1:100(4° overnight). High-pressure and temperature Tris-EDTA,pH8.0 was used for antigen retrieval. Negative contrl (right) obtaned from antibody was pre-absorbed by immunogen peptide.



Immunohistochemistry analysis of paraffin-embedded human brain tissue, using BCAR3 Antibody. The picture on the right is blocked with the synthesized peptide.



Western blot analysis of lysates from HeLa and Jurkat cells, using BCAR3 Antibody. The lane on the right is blocked with the synthesized peptide.