



MRP-L16 Polyclonal Antibody

| help in protein synthesis within the mitochondrion. Mitochondrial ribosomes (mitoribosomes) consist of a small 28S subunit and a large 39S subunit. They have an estimated 75% protein to rRNA composition compared to prokaryotic ribosomes, where this ratio is reversed. Another difference between mammalia mitoribosomes and prokaryotic ribosomes is that the latter contain a 5S rRNA. Among different species, the proteins comprising the mitoribosome differ greatless. | | |
|---|--------------------|--|
| Reactivity Human;Rat;Mouse; Applications WB;IHC;IF;ELISA Gene Name MRPL16 Protein Name 39S ribosomal protein L16 mitochondrial Immunogen The antiserum was produced against synthesized peptide derived from human MRPL16. AA range:171-220 Specificity MRP-L16 Polyclonal Antibody detects endogenous levels of MRP-L16 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB: 1/500 - 1/2000. IHC: 1/100 - 1/300. ELISA: 1/40000 IF 1:50-200 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms MRPL16; PNAS-111; 39S ribosomal protein L16; mitochondrial; L16mt; MRP-L16 Observed Band 28kD Cell Pathway Mitochondrion. Tissue Specificity Lung,Pheochromocytoma,Promyelocytic leukemia, | Catalog No | BYab-03979 |
| Applications WB;IHC;IF;ELISA Gene Name MRPL16 Protein Name 39S ribosomal protein L16 mitochondrial Immunogen The antiserum was produced against synthesized peptide derived from human MRPL16. AA range:171-220 Specificity MRP-L16 Polyclonal Antibody detects endogenous levels of MRP-L16 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB: 1/500 - 1/2000. IHC: 1/100 - 1/300. ELISA: 1/40000 IF 1:50-200 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms MRPL16; PNAS-111; 39S ribosomal protein L16; mitochondrial; L16mt; MRP-L16 Observed Band 28kD Cell Pathway Mitochondrion . Tissue Specificity Lung,Pheochromocytoma,Promyelocytic leukemia, function:Component of the large subunit of mitochondrial ribosome, similarity:Belongs to the ribosomal protein L16P family., Mammalian mitochondrial ribosomal proteins are encoded by nuclear genes a help in protein synthesis within the mitochondrion. Mitochondrial ribosomes, where this ratio is reversed. Another difference between mammalia mitoribosomes, where this ratio is reversed. Another difference between mammalia mitoribosomes is that the latter contain a 5S rRNA. Among different species, the proteins comprising the mitoribosome differ great | Isotype | IgG |
| Gene Name MRPL16 Protein Name 39S ribosomal protein L16 mitochondrial Immunogen The antiserum was produced against synthesized peptide derived from human MRPL16. AA range:171-220 Specificity MRP-L16 Polyclonal Antibody detects endogenous levels of MRP-L16 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB: 1/500 - 1/2000. IHC: 1/100 - 1/300. ELISA: 1/40000 IF 1:50-200 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms MRPL16; PNAS-111; 39S ribosomal protein L16; mitochondrial; L16mt; MRP-L16 Observed Band 28kD Cell Pathway Mitochondrion . Tissue Specificity Lung,Pheochromocytoma,Promyelocytic leukemia, function:Component of the large subunit of mitochondrial ribosomes, similarity:Belongs to the ribosomal protein L16P family., Background Mammalian mitochondrial ribosomal proteins are encoded by nuclear genes a help in protein synthesis within the mitochondrion. Mitochondrial ribosomes in synthesis within the mitochondrion compared to prokaryotic ribosomes | Reactivity | Human;Rat;Mouse; |
| Protein Name 39S ribosomal protein L16 mitochondrial | Applications | WB;IHC;IF;ELISA |
| Immunogen The antiserum was produced against synthesized peptide derived from human MRPL16. AA range:171-220 Specificity MRP-L16 Polyclonal Antibody detects endogenous levels of MRP-L16 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB: 1/500 - 1/2000. IHC: 1/100 - 1/300. ELISA: 1/40000 IF 1:50-200 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms MRPL16; PNAS-111; 39S ribosomal protein L16; mitochondrial; L16mt; MRP-L16 Observed Band 28kD Cell Pathway Mitochondrion . Tissue Specificity Lung,Pheochromocytoma,Promyelocytic leukemia, Function function:Component of the large subunit of mitochondrial ribosome, similarity:Belongs to the ribosomal protein L16P family., Background Mammalian mitochondrial ribosomal proteins are encoded by nuclear genes a help in protein synthesis within the mitochondrian Altochondrial ribosomes (mitoribosomes) consist of a small 28S subunit and a large 39S subunit. They have an estimated 75% protein to rRNA composition compared to prokaryotic ribosomes, and prokaryotic ribosomes i | Gene Name | MRPL16 |
| MRPL16. AA range:171-220 Specificity MRP-L16 Polyclonal Antibody detects endogenous levels of MRP-L16 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB: 1/500 - 1/2000. IHC: 1/100 - 1/300. ELISA: 1/40000 IF 1:50-200 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms MRPL16; PNAS-111; 39S ribosomal protein L16; mitochondrial; L16mt; MRP-L16 Observed Band 28kD Cell Pathway Mitochondrion . Tissue Specificity Lung,Pheochromocytoma,Promyelocytic leukemia, Function function:Component of the large subunit of mitochondrial ribosome, similarity:Belongs to the ribosomal protein L16P family., Mammalian mitochondrial ribosomal proteins are encoded by nuclear genes a help in protein synthesis within the mitochondrion. Mitochondrial ribosomes (mitoribosomes) consist of a small 28S subunit and a large 39S subunit. They have an estimated 75% protein to rRNA composition compared to prokaryotic ribosomes, where this ratio is reversed. Another difference between mammalia mitoribosomes and prokaryotic ribosomes is that the latter contain a 5S rRNA. Among different species, the proteins comprising the mitoribosome differ greati | Protein Name | 39S ribosomal protein L16 mitochondrial |
| Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB: 1/500 - 1/2000. IHC: 1/100 - 1/300. ELISA: 1/40000 IF 1:50-200 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms MRPL16; PNAS-111; 39S ribosomal protein L16; mitochondrial; L16mt; MRP-L16 Observed Band 28kD Cell Pathway Mitochondrion. Tissue Specificity Lung,Pheochromocytoma,Promyelocytic leukemia, Function function:Component of the large subunit of mitochondrial ribosome, similarity:Belongs to the ribosomal protein L16P family., Mammalian mitochondrial ribosomal proteins are encoded by nuclear genes a help in protein synthesis within the mitochondrion. Mitochondrial ribosomes (mitoribosomes) consist of a small 28S subunit and a large 39S subunit. They have an estimated 75% protein to rRNA composition compared to prokaryotic ribosomes, where this ratio is reversed. Another difference between mammalia mitoribosomes and prokaryotic ribosomes is that the latter contain a 5S rRNA. Among different species, the proteins comprising the mitoribosome differ greati | Immunogen | |
| Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB: 1/500 - 1/2000. IHC: 1/100 - 1/300. ELISA: 1/40000 IF 1:50-200 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms MRPL16; PNAS-111; 39S ribosomal protein L16; mitochondrial; L16mt; MRP-L16 Observed Band 28kD Cell Pathway Mitochondrion . Tissue Specificity Lung,Pheochromocytoma,Promyelocytic leukemia, Function function:Component of the large subunit of mitochondrial ribosome, similarity:Belongs to the ribosomal protein L16P family., Background Mammalian mitochondrial ribosomal proteins are encoded by nuclear genes a help in protein synthesis within the mitochondrion. Mitochondrial ribosomes (mitoribosomes) consist of a small 28S subunit and a large 39S subunit. They have an estimated 75% protein to rRNA composition compared to prokaryotic ribosomes, where this ratio is reversed. Another difference between mammalia mitoribosomes, where this ratio is reversed. Another difference between mammalia mitoribosomes and prokaryotic ribosomes is that the latter contain a 5S rRNA. Among different species, the proteins comprising the mitoribosome differ great | Specificity | MRP-L16 Polyclonal Antibody detects endogenous levels of MRP-L16 protein. |
| Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB: 1/500 - 1/2000. IHC: 1/100 - 1/300. ELISA: 1/40000 IF 1:50-200 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms MRPL16; PNAS-111; 39S ribosomal protein L16; mitochondrial; L16mt; MRP-L16 Observed Band 28kD Cell Pathway Mitochondrion. Tissue Specificity Lung,Pheochromocytoma,Promyelocytic leukemia, Function function:Component of the large subunit of mitochondrial ribosome.,similarity:Belongs to the ribosomal protein L16P family., Background Mammalian mitochondrial ribosomal proteins are encoded by nuclear genes a help in protein synthesis within the mitochondrion. Mitochondrial ribosomes (mitoribosomes) consist of a small 28S subunit and a large 39S subunit. They have an estimated 75% protein to rRNA compared to prokaryotic ribosomes, where this ratio is reversed. Another difference between mammalia mitoribosomes and prokaryotic ribosomes is that the latter contain a 5S rRNA. Among different species, the proteins comprising the mitoribosome differ greati | Formulation | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |
| affinity-chromatography using epitope-specific immunogen. WB: 1/500 - 1/2000. IHC: 1/100 - 1/300. ELISA: 1/40000 IF 1:50-200 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms MRPL16; PNAS-111; 39S ribosomal protein L16; mitochondrial; L16mt; MRP-L16 Observed Band 28kD Cell Pathway Mitochondrion. Tissue Specificity Lung,Pheochromocytoma,Promyelocytic leukemia, Function function:Component of the large subunit of mitochondrial ribosome.,similarity:Belongs to the ribosomal protein L16P family., Background Mammalian mitochondrial ribosomal proteins are encoded by nuclear genes a help in protein synthesis within the mitochondrion. Mitochondrial ribosomes (mitoribosomes) consist of a small 28S subunit and a large 39S subunit. They have an estimated 75% protein to rRNA composition compared to prokaryotic ribosomes, where this ratio is reversed. Another difference between mammalia mitoribosomes and prokaryotic ribosomes and prokaryotic ribosomes is that the latter contain a 5S rRNA. Among different species, the proteins comprising the mitoribosome differ greati | Source | Polyclonal, Rabbit,IgG |
| Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms MRPL16; PNAS-111; 39S ribosomal protein L16; mitochondrial; L16mt; MRP-L16 Observed Band 28kD Cell Pathway Mitochondrion . Tissue Specificity Lung,Pheochromocytoma,Promyelocytic leukemia, Function function:Component of the large subunit of mitochondrial ribosome.,similarity:Belongs to the ribosomal protein L16P family., Background Mammalian mitochondrial ribosomal proteins are encoded by nuclear genes a help in protein synthesis within the mitochondrion. Mitochondrial ribosomes (mitoribosomes) consist of a small 28S subunit and a large 39S subunit. They have an estimated 75% protein to rRNA composition compared to prokaryotic ribosomes, where this ratio is reversed. Another difference between mammalia mitoribosomes and prokaryotic ribosomes is that the latter contain a 5S rRNA. Among different species, the proteins comprising the mitoribosome differ greating th | Purification | · |
| Purity ≥90% Storage Stability -20°C/1 year Synonyms MRPL16; PNAS-111; 39S ribosomal protein L16; mitochondrial; L16mt; MRP-L16 Observed Band 28kD Cell Pathway Mitochondrion . Tissue Specificity Lung,Pheochromocytoma,Promyelocytic leukemia, Function function:Component of the large subunit of mitochondrial ribosome.,similarity:Belongs to the ribosomal protein L16P family., Background Mammalian mitochondrial ribosomal proteins are encoded by nuclear genes a help in protein synthesis within the mitochondrion. Mitochondrial ribosomes (mitoribosomes) consist of a small 28S subunit and a large 39S subunit. They have an estimated 75% protein to rRNA composition compared to prokaryotic ribosomes, where this ratio is reversed. Another difference between mammalia mitoribosomes and prokaryotic ribosomes is that the latter contain a 5S rRNA. Among different species, the proteins comprising the mitoribosome differ greatly | Dilution | WB: 1/500 - 1/2000. IHC: 1/100 - 1/300. ELISA: 1/40000 IF 1:50-200 |
| Synonyms MRPL16; PNAS-111; 39S ribosomal protein L16; mitochondrial; L16mt; MRP-L16 Observed Band 28kD Cell Pathway Mitochondrion. Lung,Pheochromocytoma,Promyelocytic leukemia, Function function:Component of the large subunit of mitochondrial ribosome.,similarity:Belongs to the ribosomal protein L16P family., Mammalian mitochondrial ribosomal proteins are encoded by nuclear genes a help in protein synthesis within the mitochondrion. Mitochondrial ribosomes (mitoribosomes) consist of a small 28S subunit and a large 39S subunit. They have an estimated 75% protein to rRNA composition compared to prokaryotic ribosomes, where this ratio is reversed. Another difference between mammalia mitoribosomes and prokaryotic ribosomes is that the latter contain a 5S rRNA. Among different species, the proteins comprising the mitoribosome differ greatly | Concentration | 1 mg/ml |
| Synonyms MRPL16; PNAS-111; 39S ribosomal protein L16; mitochondrial; L16mt; MRP-L16 Observed Band 28kD Cell Pathway Mitochondrion . Tissue Specificity Lung,Pheochromocytoma,Promyelocytic leukemia, Function function:Component of the large subunit of mitochondrial ribosome.,similarity:Belongs to the ribosomal protein L16P family., Background Mammalian mitochondrial ribosomal proteins are encoded by nuclear genes a help in protein synthesis within the mitochondrion. Mitochondrial ribosomes (mitoribosomes) consist of a small 28S subunit and a large 39S subunit. They have an estimated 75% protein to rRNA composition compared to prokaryotic ribosomes, where this ratio is reversed. Another difference between mammalia mitoribosomes and prokaryotic ribosomes is that the latter contain a 5S rRNA. Among different species, the proteins comprising the mitoribosome differ great | Purity | ≥90% |
| MRP-L16 Observed Band Z8kD Mitochondrion . Tissue Specificity Lung,Pheochromocytoma,Promyelocytic leukemia, function:Component of the large subunit of mitochondrial ribosome.,similarity:Belongs to the ribosomal protein L16P family., Mammalian mitochondrial ribosomal proteins are encoded by nuclear genes a help in protein synthesis within the mitochondrion. Mitochondrial ribosomes (mitoribosomes) consist of a small 28S subunit and a large 39S subunit. They have an estimated 75% protein to rRNA composition compared to prokaryotic ribosomes, where this ratio is reversed. Another difference between mammalia mitoribosomes and prokaryotic ribosomes is that the latter contain a 5S rRNA. Among different species, the proteins comprising the mitoribosome differ greatly | Storage Stability | -20°C/1 year |
| Cell Pathway Mitochondrion . Lung,Pheochromocytoma,Promyelocytic leukemia, function:Component of the large subunit of mitochondrial ribosome.,similarity:Belongs to the ribosomal protein L16P family., Mammalian mitochondrial ribosomal proteins are encoded by nuclear genes a help in protein synthesis within the mitochondrion. Mitochondrial ribosomes (mitoribosomes) consist of a small 28S subunit and a large 39S subunit. They have an estimated 75% protein to rRNA composition compared to prokaryotic ribosomes, where this ratio is reversed. Another difference between mammalia mitoribosomes and prokaryotic ribosomes is that the latter contain a 5S rRNA. Among different species, the proteins comprising the mitoribosome differ greatly | Synonyms | • |
| Tissue Specificity Lung,Pheochromocytoma,Promyelocytic leukemia, function:Component of the large subunit of mitochondrial ribosome.,similarity:Belongs to the ribosomal protein L16P family., Mammalian mitochondrial ribosomal proteins are encoded by nuclear genes a help in protein synthesis within the mitochondrion. Mitochondrial ribosomes (mitoribosomes) consist of a small 28S subunit and a large 39S subunit. They have an estimated 75% protein to rRNA composition compared to prokaryotic ribosomes, where this ratio is reversed. Another difference between mammalia mitoribosomes and prokaryotic ribosomes is that the latter contain a 5S rRNA. Among different species, the proteins comprising the mitoribosome differ greatly | Observed Band | 28kD |
| Function function:Component of the large subunit of mitochondrial ribosome.,similarity:Belongs to the ribosomal protein L16P family., Mammalian mitochondrial ribosomal proteins are encoded by nuclear genes a help in protein synthesis within the mitochondrion. Mitochondrial ribosomes (mitoribosomes) consist of a small 28S subunit and a large 39S subunit. They have an estimated 75% protein to rRNA composition compared to prokaryotic ribosomes, where this ratio is reversed. Another difference between mammalia mitoribosomes and prokaryotic ribosomes is that the latter contain a 5S rRNA. Among different species, the proteins comprising the mitoribosome differ greatless. | Cell Pathway | Mitochondrion . |
| Background Mammalian mitochondrial ribosomal proteins are encoded by nuclear genes a help in protein synthesis within the mitochondrion. Mitochondrial ribosomes (mitoribosomes) consist of a small 28S subunit and a large 39S subunit. They have an estimated 75% protein to rRNA composition compared to prokaryotic ribosomes, where this ratio is reversed. Another difference between mammalia mitoribosomes and prokaryotic ribosomes is that the latter contain a 5S rRNA. Among different species, the proteins comprising the mitoribosome differ greatless. | Tissue Specificity | Lung,Pheochromocytoma,Promyelocytic leukemia, |
| help in protein synthesis within the mitochondrion. Mitochondrial ribosomes (mitoribosomes) consist of a small 28S subunit and a large 39S subunit. They have an estimated 75% protein to rRNA composition compared to prokaryotic ribosomes, where this ratio is reversed. Another difference between mammalia mitoribosomes and prokaryotic ribosomes is that the latter contain a 5S rRNA. Among different species, the proteins comprising the mitoribosome differ greatless. | Function | |
| | Background | (mitoribosomes) consist of a small 28S subunit and a large 39S subunit. They have an estimated 75% protein to rRNA composition compared to prokaryotic ribosomes, where this ratio is reversed. Another difference between mammalian |

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658



国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询




| | recognition by sequence homology. This gene encodes a 39S subunit protein. [provided by RefSeq, Jul 2008], |
|---------------------------|---|
| matters needing attention | Avoid repeated freezing and thawing! |
| Usage suggestions | This product can be used in immunological reaction related experiments. For more information, please consult technical personnel. |







Products Images



Western Blot analysis of various cells using MRP-L16 Polyclonal Antibody diluted at 1:2000



Immunohistochemistry analysis of paraffin-embedded human heart tissue, using MRPL16 Antibody. The picture on the right is blocked with the synthesized peptide.



Western blot analysis of lysates from 293 cells, using MRPL16 Antibody. The lane on the right is blocked with the synthesized peptide.

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658

国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询



