



RGS7 Polyclonal Antibody

Catalog No	BYab-04125
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB;ELISA;IHC
Gene Name	RGS7
Protein Name	Regulator of G-protein signaling 7
Immunogen	The antiserum was produced against synthesized peptide derived from human RGS7. AA range:155-204
Specificity	RGS7 Polyclonal Antibody detects endogenous levels of RGS7 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000;IHC-p 1:50-300; ELISA 2000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	RGS7; Regulator of G-protein signaling 7; RGS7
Observed Band	65kD
Cell Pathway	Cytoplasm, cytosol . Cytoplasm . Cell membrane . Membrane ; Peripheral membrane protein ; Cytoplasmic side . Interaction with PKD1 promotes location at the cell membrane (PubMed:10339594). Interaction with RGS7BP promotes location at the cell membrane (PubMed:15897264). .
Tissue Specificity	Brain,
Function	function:Inhibits signal transduction by increasing the GTPase activity of G protein alpha subunits thereby driving them into their inactive GDP-bound form. Activity on G(o)-alpha is specifically enhanced by the RGS6/GNG5 dimer. May play a role in synaptic vesicle exocytosis. May play important role in the rapid regulation of neuronal excitability and the cellular responses to short-lived stimulations.,PTM:Palmitoylated.,PTM:Phosphorylation and subsequent interaction with 14-3-3 proteins inhibits GAP activity.,similarity:Contains 1 DEP domain.,similarity:Contains 1 G protein gamma domain.,similarity:Contains 1 RGS domain.,subunit:Heterodimer with GNG5. Interacts with RGS7BP, leading to

Nanjing BYabscience technology Co.,Ltd



regulate the subcellular location of the heterodimer formed with Gbeta5 (By similarity). Interacts with 14-3-3 protein Tau and SNAP25BP.,

Background

function:Inhibits signal transduction by increasing the GTPase activity of G protein alpha subunits thereby driving them into their inactive GDP-bound form. Activity on G(o)-alpha is specifically enhanced by the RGS6/GNG5 dimer. May play a role in synaptic vesicle exocytosis. May play important role in the rapid regulation of neuronal excitability and the cellular responses to short-lived stimulations.,PTM:Palmitoylated.,PTM:Phosphorylation and subsequent interaction with 14-3-3 proteins inhibits GAP activity.,similarity:Contains 1 DEP domain.,similarity:Contains 1 G protein gamma domain.,similarity:Contains 1 RGS domain.,subunit:Heterodimer with GNG5. Interacts with RGS7BP, leading to regulate the subcellular location of the heterodimer formed with Gbeta5 (By similarity). Interacts with 14-3-3 protein Tau and SNAP25BP.,

matters needing attention


Avoid repeated freezing and thawing!

Usage suggestions


This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.



Products Images



Western blot analysis of lysates from Jurkat cells, using RGS7 Antibody. The lane on the right is blocked with the synthesized peptide.



Immunohistochemical analysis of paraffin-embedded human Gastric adenocarcinoma. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).