



# Chymase Polyclonal Antibody

<b>Catalog No</b>	BYab-04304
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Rat;Mouse;
<b>Applications</b>	WB;ELISA
<b>Gene Name</b>	CMA1
<b>Protein Name</b>	Chymase
<b>Immunogen</b>	Synthesized peptide derived from the Internal region of human Chymase.
<b>Specificity</b>	Chymase Polyclonal Antibody detects endogenous levels of Chymase protein.
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source</b>	Polyclonal, Rabbit,IgG
<b>Purification</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Dilution</b>	Western Blot: 1/500 - 1/2000. ELISA: 1/20000. Not yet tested in other applications.
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	CMA1; CYH; CYM; Chymase; Alpha-chymase; Mast cell protease I
<b>Observed Band</b>	27kD
<b>Cell Pathway</b>	Secreted. Cytoplasmic granule. Mast cell granules.
<b>Tissue Specificity</b>	Mast cells in lung, heart, skin and placenta. Expressed in both normal skin and in urticaria pigmentosa lesions.
<b>Function</b>	catalytic activity:Preferential cleavage: Phe- -Xaa > Tyr- -Xaa > Trp- -Xaa > Leu- -Xaa.,function:Major secreted protease of mast cells with suspected roles in vasoactive peptide generation, extracellular matrix degradation, and regulation of gland secretion.,similarity:Belongs to the peptidase S1 family.,similarity:Belongs to the peptidase S1 family. Granzyme subfamily.,similarity:Contains 1 peptidase S1 domain.,subcellular location:Mast cell granules.,tissue specificity:Mast cells in lung, heart, skin and placenta.,
<b>Background</b>	chymase 1(CMA1) Homo sapiens This gene encodes a chymotryptic serine proteinase that belongs to the peptidase family S1. It is expressed in mast cells and is thought to function in the degradation of the extracellular matrix, the regulation of submucosal gland secretion, and the generation of vasoactive

Nanjing BYabscience technology Co.,Ltd



peptides. In the heart and blood vessels, this protein, rather than angiotensin converting enzyme, is largely responsible for converting angiotensin I to the vasoactive peptide angiotensin II. Alternative splicing results in multiple variants. [provided by RefSeq, Apr 2015],


**matters needing attention**

Avoid repeated freezing and thawing!

**Usage suggestions**

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## Products Images



Western Blot analysis of mouse lung cells using Chymase Polyclonal Antibody. Secondary antibody(catalog#:RS0002) was diluted at 1:20000