



Noggin mouse mAb

Catalog No	BYab-04479
Isotype	IgG
Reactivity	0
Applications	WB;ICC
Gene Name	nog
Protein Name	
Immunogen	Purified recombinant human Noggin protein fragments expressed in E.coli.
Specificity	Transfected Only.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse
Purification	The antibody was affinity-purified from mouse ascites by affinity-chromatography using epitope-specific immunogen.
Dilution	wb 1:1000 icc 1:100
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	NOG;NOGG_HUMAN;Noggin;SYM 1;SYM1;Symphalangism 1 (proximal);Synostoses (multiple) syndrome 1;SYNS 1;SYNS1.
Observed Band	26kD
Cell Pathway	Secreted.
Tissue Specificity	Placenta,Prostate,Temporal cortex,
Function	disease:Defects in NOG are a cause of stapes ankylosis with broad thumb and toes [MIM:184460]. Stapes ankylosis with broad thumb and toes is a congenital autosomal dominant disorder that includes hyperopia, a hemicylindrical nose, broad thumbs, great toes, and other minor skeletal anomalies but lacked carpal and tarsal fusion and symphalangism.,disease:Defects in NOG are a cause of symphalangism proximal syndrome (SYM1) [MIM:185800]. SYM1 is characterized by the hereditary absence of the proximal interphalangeal (PIP) joints (Cushing symphalangism). Severity of PIP joint involvement diminishes towards the radial side. Distal interphalangeal joints are less frequently involved and metacarpophalangeal joints are rarely affected whereas carpal bone malformation and fusion are common. In the lower extremities, tarsal bone

Nanjing BYabscience technology Co.,Ltd



coalition is common. Conductive hearing loss is seen and is due to fusi

Background

The secreted polypeptide, encoded by this gene, binds and inactivates members of the transforming growth factor-beta (TGF-beta) superfamily signaling proteins, such as bone morphogenetic protein-4 (BMP4). By diffusing through extracellular matrices more efficiently than members of the TGF-beta superfamily, this protein may have a principal role in creating morphogenic gradients. The protein appears to have pleiotropic effect, both early in development as well as in later stages. It was originally isolated from *Xenopus* based on its ability to restore normal dorsal-ventral body axis in embryos that had been artificially ventralized by UV treatment. The results of the mouse knockout of the ortholog suggest that it is involved in numerous developmental processes, such as neural tube fusion and joint formation. Recently, several dominant human NOG mutations in unrelated families with proximal symphalangism (SYM1) and mu

matters needing attention

Avoid repeated freezing and thawing!


Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.


Nanjing BYabscience technology Co.,Ltd



Products Images



Western blot detection of Noggin in CHO-K1 cell lysates over-expressing Noggin-PDGFR transmembrane domain fused protein using Noggin mouse mAb (1:1000 diluted). Predicted band size: 26KDa. Observed band size: 37KDa.



Immunocytochemistry staining of HeLa cells surface-expressing Noggin-PDGFR transmembrane domain fused protein using Noggin mouse mAb (dilution 1:100).