




PDE3A Polyclonal Antibody

Catalog No BYab-04938 Isotype IgG Reactivity Human;Mouse;Rat Applications WB;ELISA Gene Name PDE3A Protein Name cGMP-inhibited 3',5'-cyclic phosphodiesterase A (EC 3.1.4.17) (Cyclic GMP-inhibited phosphodiesterase A) (CGI-PDE A) Immunogen Synthesized peptide derived from human protein . at AA range: 250-330 Specificity PDE3A Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Cell Pathway Observed Band 125kD Cell Pathway Membrane; Multi-pass membrane protein . Cytoplasm, cytosol . Tissue Specificity Blood,Epithelium,Liver,Myocardium,Penis,Platelet, Function catalytic activity,Nucleoside 3',5'-cyclic phosphate + H(2)O = nucleoside 5			
Reactivity Human; Mouse; Rat Applications WB; ELISA Gene Name PDE3A Protein Name cGMP-inhibited 3',5'-cyclic phosphodiesterase A (CGI-PDE A) Immunogen Synthesized peptide derived from human protein . at AA range: 250-330 Specificity PDE3A Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 125kD Cell Pathway Membrane; Multi-pass membrane protein . Cytoplasm, cytosol . Tissue Specificity Blood, Epithelium, Liver, Myocardium, Penis, Platelet, Function catalytic activity. Nucleoside 3', 5'-cyclic phosphate + H(2)O = nucleoside 5'-phosphate, enzyme regulation; Inhibited of yo GMP, incrition: Hydrolyzes both cyclic AMP (CAMP) and cyclic GMP (GAMP), online information; PDE3 entry, similarity: Belongs to the cyclic hucleotide phosphodiesterase (GI-PDE) family. GGI	Catalog No	BYab-04938	
Applications WB;ELISA Gene Name PDE3A Protein Name cGMP-inhibited 31,51-cyclic phosphodiesterase A (EC 3.1.4.17) (Cyclic GMP-inhibited phosphodiesterase A) (CGI-PDE A) Immunogen Synthesized peptide derived from human protein . at AA range: 250-330 Specificity PDE3A Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 125kD Cell Pathway Membrane; Multi-pass membrane protein . Cytoplasm, cytosol . Tissue Specificity Blood, Epithelium, Liver, Myocardium, Penis, Platelet, Function catalytic activity: Nucleoside 3: 51-cyclic phosphate + H(2)O = nucleoside 51-phosphate, enzyme regulation: Inhibited by cGMP, function PDE3 entry, similarity: Belongs to the cyclic nucleotide phosphodiesterase family. This gene encodes a member of the cGMP-inhibited cyclic nucleotide phosphodiesterase (GI-PDE) family. CGI-PDE parzymes hydyoze both cAMP and CGMP, and play critical roles in many cellular processes by regulating the amplitude and duration of intracellular cyclic nucleotide in cardiovascular function by regulating vascular smooth muscle contraction and	Isotype	IgG	
Gene Name PDE3A Protein Name cGMP-inhibited 3',5'-cyclic phosphodiesterase A (EC 3.1.4.17) (Cyclic GMP-inhibited phosphodiesterase A) (CGI-PDE A) Immunogen Synthesized peptide derived from human protein . at AA range: 250-330 Specificity PDE3A Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 125kD Cell Pathway Membrane; Multi-pass membrane protein . Cytoplasm, cytosol . Tissue Specificity Blood,Epithelium,Liver,Myocardium,Penis,Platelet, Function catalytic activity:Nucleoside 3',5'-cyclic phosphate + H(2)O = nucleoside 5'-phosphate, enzyme regulation:Inhibited by cGMP, function:Hydrolyzes both cyclic AMP (cAMP) and cyclic GMP (cGMP), online information:PDE3 entry, similarity. Belongs to the cyclic nucleotide phosphodiesterase family. Background This gene encodes a member of the cGMP-inhibited cyclic nucleo	Reactivity	Human;Mouse;Rat	
Protein Name	Applications	WB;ELISA	
Immunogen Synthesized peptide derived from human protein . at AA range: 250-330 Specificity PDE3A Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 125kD Cell Pathway Membrane; Multi-pass membrane protein. Cytoplasm, cytosol. Tissue Specificity Blood,Epithelium,Liver,Myocardium,Penis,Platelet, Function catalytic activity.Nucleoside 3',5'-cyclic phosphate + H(2)O = nucleoside 5'-phosphate.,enzyme regulation:Inhibited by cGMP, function:Hydrolyzes both cyclic AMP (cAMP) and cyclic GMP (cGMP), online information:PDE3 entry, similarity.Belongs to the cyclic nucleotide phosphodiesterase family. Background This gene encodes a member of the cGMP-inhibited cyclic nucleotide phosphodiesterase (cGI-PDE) family. cGI-PDE enzymes hydrolyze both cAMP and CGMP, and play critical roles in many cellular processes by regulating the amplitude and duration of intracellular cyclic nucleotide signals. The encoded protein mediates pla	Gene Name	PDE3A	
Specificity PDE3A Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 125kD Cell Pathway Membrane; Multi-pass membrane protein. Cytoplasm, cytosol. Tissue Specificity Blood,Epithelium,Liver,Myocardium,Penis,Platelet, Function catalytic activity:Nucleoside 3',5'-cyclic phosphate + H(2)O = nucleoside 5'-phosphate.enzyme regulation:Inhibited by cGMP, function:Hydrolyzes both cyclic AMP (cAMP) and cyclic GMP (cGMP), online information:PDE3 entry, similarity:Belongs to the cyclic nucleotide phosphodiesterase familly., Background This gene encodes a member of the cGMP-inhibited cyclic nucleotide phosphodiesterase (cGI-PDE) family. cGI-PDE enzymes hydrolyze both cAMP and cGMP, and play critical roles in many cellular processes by regulating the amplitude and duration of intracellular cyclic nucleotide signals. The encoded protein mediates platelet aggregation and also plays important roles in cardiovascular function by regulating vascular smooth muscle contraction and	Protein Name	cGMP-inhibited 3',5'-cyclic phosphodiesterase A (EC 3.1.4.17) (Cyclic GMP-inhibited phosphodiesterase A) (CGI-PDE A)	
Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 125kD Cell Pathway Membrane; Multi-pass membrane protein. Cytoplasm, cytosol. Tissue Specificity Blood,Epithelium,Liver,Myocardium,Penis,Platelet, Function catalytic activity:Nucleoside 3',5'-cyclic phosphate + H(2)O = nucleoside 5'-phosphate, enzyme regulation:Inhibited by cGMP, function:Hydrolyzes both cyclic AMP (cAMP) and cyclic GMP (cGMP),online information:PDE3 entry,similarity:Belongs to the cyclic nucleotide phosphodiesterase family. Background This gene encodes a member of the cGMP-inhibited cyclic nucleotide phosphodiesterase (cGH-PDE) family. cGI-PDE enzymes hydrolyze both cAMP and cGMP, and play critical roles in many cellular processes by regulating the amplitude and duration of intracellular cyclic nucleotide signals. The encoded protein mediates platelet aggregation and also plays important roles in cardiovascular function by regulating vascular smooth muscle contraction and	Immunogen	Synthesized peptide derived from human protein . at AA range: 250-330	
Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 125kD Cell Pathway Membrane ; Multi-pass membrane protein . Cytoplasm, cytosol . Tissue Specificity Blood,Epithelium,Liver,Myocardium,Penis,Platelet, Function catalytic activity:Nucleoside 3',5'-cyclic phosphate + H(2)O = nucleoside 5'-phosphate,enzyme regulation:Inhibited by cGMP, function:Hydrolyzes both cyclic AMP (cAMP) and cyclic GMP (cGMP), online information:PDE3 entry, similanity:Belongs to the cyclic nucleotide phosphodiesterase family. This gene encodes a member of the cGMP-inhibited cyclic nucleotide phosphodiesterase (cGI-PDE) family. cGI-PDE enzymes hydrolyze both cAMP and cGMP, and play critical roles in many cellular processes by regulating the amplitude and duration of intracellular cyclic nucleotide signals. The encoded protein mediates platelet aggregation and also plays important roles in cardiovascular function by regulating vascular smooth muscle contraction and	Specificity	PDE3A Polyclonal Antibody detects endogenous levels of protein.	
Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 125kD Cell Pathway Membrane ; Multi-pass membrane protein . Cytoplasm, cytosol . Tissue Specificity Blood,Epithelium,Liver,Myocardium,Penis,Platelet, Function catalytic activity:Nucleoside 3',5'-cyclic phosphate + H(2)O = nucleoside 5'-phosphate.enzyme regulation:Inhibited by cGMP_function:Hydrolyzes both cyclic AMP (cAMP) and cyclic GMP (cGMP), innetion:Hydrolyzes both cyclic AMP (cAMP) and cyclic GMP (cGMP)-inhibited cyclic nucleotide phosphodiesterase family., Background This gene encodes a member of the cGMP-inhibited cyclic nucleotide phosphodiesterase (cGI-PDE) family. cGI-PDE enzymes hydrolyze both cAMP and cGMP, and play critical roles in many cellular processes by regulating the amplitude and duration of intracellular cyclic nucleotide signals. The encoded protein mediates platelet aggregation and also plays important roles in cardiovascular function by regulating yascular smooth muscle contraction and	Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.	
affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 125kD Cell Pathway Membrane ; Multi-pass membrane protein . Cytoplasm, cytosol . Tissue Specificity Blood,Epithelium,Liver,Myocardium,Penis,Platelet, Function catalytic activity:Nucleoside 3',5'-cyclic phosphate + H(2)O = nucleoside 5'-phosphate,enzyme regulation:Inhibited by cGMP_intinction:Hydrolyzes both cyclic AMP (aCMP) and cyclic GMP (aCMP),online information:PDE3 entry,similarity:Belongs to the cyclic nucleotide phosphodiesterase family., Background This gene encodes a member of the cGMP-inhibited cyclic nucleotide phosphodiesterase (cGI-PDE) family. cGI-PDE enzymes hydrolyze both cAMP and cGMP, and play critical roles in many cellular processes by regulating the amplitude and duration of intracellular cyclic nucleotide signals. The encoded protein mediates platelet aggregation and also plays important roles in cardiovascular function by regulating vascular smooth muscle contraction and	Source	Polyclonal, Rabbit,IgG	
Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 125kD Cell Pathway Membrane; Multi-pass membrane protein. Cytoplasm, cytosol. Tissue Specificity Blood,Epithelium,Liver,Myocardium,Penis,Platelet, Function catalytic activity:Nucleoside 3',5'-cyclic phosphate + H(2)O = nucleoside 5'-phosphate.enzyme regulation:Inhibited by cGMP.function:Hydrolyzes both cyclic AMP (cAMP) and cyclic GMP (cGMP).online information:PDE3 entry,similarity:Belongs to the cyclic nucleotide phosphodiesterase family., Background This gene encodes a member of the cGMP-inhibited cyclic nucleotide phosphodiesterase (cGI-PDE) family. cGI-PDE enzymes hydrolyze both cAMP and cGMP, and play critical roles in many cellular processes by regulating the amplitude and duration of intracellular cyclic nucleotide signals. The encoded protein mediates platelet aggregation and also plays important roles in cardiovascular function by regulating vascular smooth muscle contraction and	Purification		
Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 125kD Cell Pathway Membrane; Multi-pass membrane protein. Cytoplasm, cytosol. Tissue Specificity Blood,Epithelium,Liver,Myocardium,Penis,Platelet, Function catalytic activity:Nucleoside 3',5'-cyclic phosphate + H(2)O = nucleoside 5'-phosphate.,enzyme regulation:Inhibited by cGMP.,function:Hydrolyzes both cyclic AMP (cAMP) and cyclic GMP (cGMP).,online information:PDE3 entry,similarity:Belongs to the cyclic nucleotide phosphodiesterase family., This gene encodes a member of the cGMP-inhibited cyclic nucleotide phosphodiesterase (cGI-PDE) family. cGI-PDE enzymes hydrolyze both cAMP and cGMP, and play critical roles in many cellular processes by regulating the amplitude and duration of intracellular cyclic nucleotide signals. The encoded protein mediates platelet aggregation and also plays important roles in cardiovascular function by regulating vascular smooth muscle contraction and	Dilution	WB 1:500-2000 ELISA 1:5000-20000	
Storage Stability -20°C/1 year Synonyms Observed Band 125kD Cell Pathway Membrane; Multi-pass membrane protein. Cytoplasm, cytosol. Tissue Specificity Blood,Epithelium,Liver,Myocardium,Penis,Platelet, Function catalytic activity:Nucleoside 3',5'-cyclic phosphate + H(2)O = nucleoside 5'-phosphate.,enzyme regulation:Inhibited by cGMP.,function:Hydrolyzes both cyclic AMP (cAMP) and cyclic GMP (cGMP).,online information:PDE3 entry,similarity:Belongs to the cyclic nucleotide phosphodiesterase family., This gene encodes a member of the cGMP-inhibited cyclic nucleotide phosphodiesterase (cGI-PDE) family. cGI-PDE enzymes hydrolyze both cAMP and cGMP, and play critical roles in many cellular processes by regulating the amplitude and duration of intracellular cyclic nucleotide signals. The encoded protein mediates platelet aggregation and also plays important roles in cardiovascular function by regulating vascular smooth muscle contraction and	Concentration	1 mg/ml	
Synonyms Deserved Band 125kD	Purity	≥90%	
Observed Band Cell Pathway Membrane; Multi-pass membrane protein. Cytoplasm, cytosol. Blood,Epithelium,Liver,Myocardium,Penis,Platelet, Catalytic activity:Nucleoside 3',5'-cyclic phosphate + H(2)O = nucleoside 5'-phosphate.,enzyme regulation:Inhibited by cGMP.,function:Hydrolyzes both cyclic AMP (cAMP) and cyclic GMP (cGMP).,online information:PDE3 entry,similarity:Belongs to the cyclic nucleotide phosphodiesterase family. Background This gene encodes a member of the cGMP-inhibited cyclic nucleotide phosphodiesterase (cGI-PDE) family. cGI-PDE enzymes hydrolyze both cAMP and cGMP, and play critical roles in many cellular processes by regulating the amplitude and duration of intracellular cyclic nucleotide signals. The encoded protein mediates platelet aggregation and also plays important roles in cardiovascular function by regulating vascular smooth muscle contraction and	04 1 1114	-20°C/1 year	
Cell Pathway Membrane; Multi-pass membrane protein. Cytoplasm, cytosol. Blood, Epithelium, Liver, Myocardium, Penis, Platelet, catalytic activity: Nucleoside 3',5'-cyclic phosphate + H(2)O = nucleoside 5'-phosphate., enzyme regulation: Inhibited by cGMP., function: Hydrolyzes both cyclic AMP (cAMP) and cyclic GMP (cGMP)., online information: PDE3 entry, similarity: Belongs to the cyclic nucleotide phosphodiesterase family., This gene encodes a member of the cGMP-inhibited cyclic nucleotide phosphodiesterase (cGI-PDE) family. cGI-PDE enzymes hydrolyze both cAMP and cGMP, and play critical roles in many cellular processes by regulating the amplitude and duration of intracellular cyclic nucleotide signals. The encoded protein mediates platelet aggregation and also plays important roles in cardiovascular function by regulating vascular smooth muscle contraction and	Storage Stability	-20°C/1 year	
Tissue Specificity Blood,Epithelium,Liver,Myocardium,Penis,Platelet, catalytic activity:Nucleoside 3',5'-cyclic phosphate + H(2)O = nucleoside 5'-phosphate.,enzyme regulation:Inhibited by cGMP.,function:Hydrolyzes both cyclic AMP (cAMP) and cyclic GMP (cGMP).,online information:PDE3 entry,similarity:Belongs to the cyclic nucleotide phosphodiesterase family., This gene encodes a member of the cGMP-inhibited cyclic nucleotide phosphodiesterase (cGI-PDE) family. cGI-PDE enzymes hydrolyze both cAMP and cGMP, and play critical roles in many cellular processes by regulating the amplitude and duration of intracellular cyclic nucleotide signals. The encoded protein mediates platelet aggregation and also plays important roles in cardiovascular function by regulating vascular smooth muscle contraction and		-20°C/1 year	
Function catalytic activity:Nucleoside 3',5'-cyclic phosphate + H(2)O = nucleoside 5'-phosphate.,enzyme regulation:Inhibited by cGMP.,function:Hydrolyzes both cyclic AMP (cAMP) and cyclic GMP (cGMP).,online information:PDE3 entry,similarity:Belongs to the cyclic nucleotide phosphodiesterase family., This gene encodes a member of the cGMP-inhibited cyclic nucleotide phosphodiesterase (cGI-PDE) family. cGI-PDE enzymes hydrolyze both cAMP and cGMP, and play critical roles in many cellular processes by regulating the amplitude and duration of intracellular cyclic nucleotide signals. The encoded protein mediates platelet aggregation and also plays important roles in cardiovascular function by regulating vascular smooth muscle contraction and	Synonyms		
5'-phósphate.,énzyme regulation:lnhibited by cGMP.,fùnction:Hydrolyzes both cyclic AMP (cAMP) and cyclic GMP (cGMP).,online information:PDE3 entry,similarity:Belongs to the cyclic nucleotide phosphodiesterase family., This gene encodes a member of the cGMP-inhibited cyclic nucleotide phosphodiesterase (cGI-PDE) family. cGI-PDE enzymes hydrolyze both cAMP and cGMP, and play critical roles in many cellular processes by regulating the amplitude and duration of intracellular cyclic nucleotide signals. The encoded protein mediates platelet aggregation and also plays important roles in cardiovascular function by regulating vascular smooth muscle contraction and	Synonyms Observed Band	125kD	
phosphodiesterase (cGI-PDE) family. cGI-PDE enzymes hydrolyze both cAMP and cGMP, and play critical roles in many cellular processes by regulating the amplitude and duration of intracellular cyclic nucleotide signals. The encoded protein mediates platelet aggregation and also plays important roles in cardiovascular function by regulating vascular smooth muscle contraction and	Synonyms Observed Band Cell Pathway	125kD Membrane ; Multi-pass membrane protein . Cytoplasm, cytosol .	
	Synonyms Observed Band Cell Pathway Tissue Specificity	125kD Membrane; Multi-pass membrane protein. Cytoplasm, cytosol. Blood,Epithelium,Liver,Myocardium,Penis,Platelet, catalytic activity:Nucleoside 3',5'-cyclic phosphate + H(2)O = nucleoside 5'-phosphate.,enzyme regulation:Inhibited by cGMP.,function:Hydrolyzes both cyclic AMP (cAMP) and cyclic GMP (cGMP).,online information:PDE3	

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658







	congestive heart failure. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene. [provided by RefSeq, Sep 2011],
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions This product can be used in immunological reaction related experiments. F more information, please consult technical personnel.	

Products Images

Nanjing BYabscience technology Co.,Ltd