





Catalog No	BYab-05044	
Isotype	IgG	
Reactivity	Human;Rat;Mouse;	
Applications	WB;ELISA	
Gene Name	CYP2F1	
Protein Name	Cytochrome P450 2F1 (EC 1.14.14.1) (CYPIIF1)	
Immunogen	Synthesized peptide derived from human protein . at AA range: 210-290	
Specificity	CP2F1 Polyclonal Antibody detects endogenous levels of protein.	
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.	
Source	Polyclonal, Rabbit,IgG	
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.	
Dilution	WB 1:500-2000 ELISA 1:5000-20000	
Concentration	1 mg/ml	
Purity	≥90%	
Storage Stability	-20°C/1 year	
Synonyms		
Observed Band	54kD	
Cell Pathway	Endoplasmic reticulum membrane ; Peripheral membrane protein . Microsome membrane ; Peripheral membrane protein .	
Tissue Specificity	Expressed in lung. Rarely detected in liver and placenta.	
Function	catalytic activity:RH + reduced flavoprotein + O(2) = ROH + oxidized flavoprotein + H(2)O.,cofactor:Heme group.,function:Is able to dealkylate ethoxycoumarin, propoxycoumarin, and pentoxyresorufin but possesses no activity toward ethoxyresorufin and only trace dearylation activity toward benzyloxyresorufin. Bioactivates 3-methylindole (3MI) by dehydrogenation to the putative electrophile 3-methylene-indolenine.,similarity:Belongs to the cytochrome P450 family.,	
Background	This gene encodes a member of the cytochrome P450 superfamily of enzymes. The cytochrome P450 proteins are monooxygenases which catalyze many reactions involved in drug metabolism and synthesis of cholesterol, steroids and other lipids. This protein localizes to the endoplasmic reticulum and is known to dehydrogenate 3-methylindole, an endogenous toxin derived from the	

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658



国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询



	fermentation of tryptophan, as well as xenobiotic substrates such as naphthalene and ethoxycoumarin. This gene is part of a large cluster of cytochrome P450 genes from the CYP2A, CYP2B and CYP2F subfamilies on chromosome 19q. [provided by RefSeq, Jul 2008],	
matters needing attention	Avoid repeated freezing and thawing!	
Usage suggestions	e suggestions  This product can be used in immunological reaction related experiments. Fo more information, please consult technical personnel.	

Products Images

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658