



IFI44 Polyclonal Antibody

Background IFI44 family., function:This protein aggregates to form microtubular structuresinduction:By		
Reactivity Human;Rat;Mouse; Applications WB;ELISA Gene Name IFI44 MTAP44 Protein Name Interferon-induced protein 44 (p44) (Microtubule-associated protein 44) Immunogen Synthesized peptide derived from human protein . at AA range: 310-390 Specificity IFI44 Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 48kD Cell Pathway Cytoplasm. Tissue Specificity Lung, Function function:This protein aggregates to form microtubular structures, induction:By alpha and beta interferons, but not by gamma interferons, similarity:Belongs to the IFI44 familly., Movid repeated freezing and thawing!	Catalog No	BYab-05085
Applications WB;ELISA Gene Name IFI44 MTAP44 Protein Name Interferon-induced protein 44 (p44) (Microtubule-associated protein 44) Immunogen Synthesized peptide derived from human protein . at AA range: 310-390 Specificity IFI44 Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 48kD Cell Pathway Cytoplasm. Tissue Specificity Lung, Function function:This protein aggregates to form microtubular structures, induction:By alpha and beta interferons, but not by gamma interferons., similarity:Belongs to the IFI44 familly. matters needing Avoid repeated freezing and thawing!	Isotype	IgG
Gene Name IFI44 MTAP44 Protein Name Interferon-induced protein 44 (p44) (Microtubule-associated protein 44) Immunogen Synthesized peptide derived from human protein . at AA range: 310-390 Specificity IFI44 Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 48kD Cell Pathway Cytoplasm . Tissue Specificity Lung, Function function:This protein aggregates to form microtubular structures, induction:By alpha and beta interferons, but not by gamma interferons, similarity:Belongs to the IFI44 family., Matters needing Avoid repeated freezing and thawing!	Reactivity	Human;Rat;Mouse;
Protein Name Interferon-induced protein 44 (p44) (Microtubule-associated protein 44) Immunogen Synthesized peptide derived from human protein . at AA range: 310-390 Specificity IFI44 Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 48kD Cell Pathway Cytoplasm . Tissue Specificity Lung, Function function:This protein aggregates to form microtubular structures.,induction:By alpha and beta interferons, but not by gamma interferons.,similarity:Belongs to the IFI44 family., Background function:This protein aggregates to form microtubular structures.,induction:By alpha and beta interferons, but not by gamma interferons.,similarity:Belongs to the IFI44 family., More and the protein aggregates to form microtubular structures.,induction:By alpha and beta interferons, but not by gamma interferons.,similarity:Belongs to the IFI44 family., More and the protein aggregates to form microtubular structures.,induction:By alpha and beta interferons, but not by gamma interferons.,similarity:Belongs to the IFI44 family., More and the protein aggregates to form microtubular structures.,induction:By alpha and beta interferons, but not by gamma interferons.,similarity:Belongs to the IFI44 family.,	Applications	WB;ELISA
Immunogen Synthesized peptide derived from human protein . at AA range: 310-390	Gene Name	IFI44 MTAP44
Specificity IFI44 Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 48kD Cell Pathway Cytoplasm Lung, Function function:This protein aggregates to form microtubular structures, induction:By alpha and beta interferons, but not by gamma interferons, similarity:Belongs to the IFI44 family. Background function:This protein aggregates to form microtubular structures, induction:By alpha and beta interferons, but not by gamma interferons, similarity:Belongs to the IFI44 family. Matters needing Avoid repeated freezing and thawing!	Protein Name	Interferon-induced protein 44 (p44) (Microtubule-associated protein 44)
Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 48kD Cell Pathway Cytoplasm . Tissue Specificity Lung, Function function:This protein aggregates to form microtubular structures, induction:By alpha and beta interferons, but not by gamma interferons.,similarity:Belongs to the IFI44 family., Background function:This protein aggregates to form microtubular structures, induction:By alpha and beta interferons, but not by gamma interferons.,similarity:Belongs to the IFI44 family.	Immunogen	Synthesized peptide derived from human protein . at AA range: 310-390
Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 48kD Cell Pathway Cytoplasm . Tissue Specificity Lung, Function function:This protein aggregates to form microtubular structures.,induction:By alpha and beta interferons, but not by gamma interferons.,similarity:Belongs to the IFI44 family., Background function:This protein aggregates to form microtubular structures.,induction:By alpha and beta interferons, but not by gamma interferons.,similarity:Belongs to the IFI44 family. matters needing Avoid repeated freezing and thawing!	Specificity	IFI44 Polyclonal Antibody detects endogenous levels of protein.
Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 48kD Cell Pathway Cytoplasm. Tissue Specificity Lung, Function function:This protein aggregates to form microtubular structures.,induction:By alpha and beta interferons, but not by gamma interferons.,similarity:Belongs to the IFI44 familly., Background function:This protein aggregates to form microtubular structures.,induction:By alpha and beta interferons, but not by gamma interferons.,similarity:Belongs to the IFI44 familly., Matters needing Avoid repeated freezing and thawing!	Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 48kD Cell Pathway Cytoplasm . Tissue Specificity Lung, Function function:This protein aggregates to form microtubular structures.,induction:By alpha and beta interferons, but not by gamma interferons.,similarity:Belongs to the IFI44 family., Background function:This protein aggregates to form microtubular structures.,induction:By alpha and beta interferons, but not by gamma interferons.,similarity:Belongs to the IFI44 family., Matters needing Avoid repeated freezing and thawing!	Source	Polyclonal, Rabbit,IgG
Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 48kD Cell Pathway Cytoplasm . Tissue Specificity Lung, Function function:This protein aggregates to form microtubular structures.,induction:By alpha and beta interferons, but not by gamma interferons.,similarity:Belongs to the IF144 family., Background function:This protein aggregates to form microtubular structures.,induction:By alpha and beta interferons, but not by gamma interferons.,similarity:Belongs to the IF144 family., matters needing Avoid repeated freezing and thawing!	Purification	
Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 48kD Cell Pathway Cytoplasm . Tissue Specificity Lung, Function function:This protein aggregates to form microtubular structuresinduction:By alpha and beta interferons, but not by gamma interferonssimilarity:Belongs to the IFI44 family Background function:This protein aggregates to form microtubular structuresinduction:By alpha and beta interferons, but not by gamma interferonssimilarity:Belongs to the IFI44 family Matters needing Avoid repeated freezing and thawing!	Dilution	WB 1:500-2000 ELISA 1:5000-20000
Storage Stability -20°C/1 year Synonyms Observed Band 48kD Cell Pathway Cytoplasm. Tissue Specificity Lung, Function function:This protein aggregates to form microtubular structures.,induction:By alpha and beta interferons, but not by gamma interferons.,similarity:Belongs to the IFI44 family., Background function:This protein aggregates to form microtubular structures.,induction:By alpha and beta interferons, but not by gamma interferons.,similarity:Belongs to the IFI44 family., matters needing Avoid repeated freezing and thawing!	Concentration	1 mg/ml
Synonyms Observed Band 48kD Cell Pathway Cytoplasm . Tissue Specificity Lung, Function function:This protein aggregates to form microtubular structures.,induction:By alpha and beta interferons, but not by gamma interferons.,similarity:Belongs to the IF144 family., Background function:This protein aggregates to form microtubular structures.,induction:By alpha and beta interferons, but not by gamma interferons.,similarity:Belongs to the IF144 family., matters needing Avoid repeated freezing and thawing!	Purity	≥90%
Observed Band Cell Pathway Cytoplasm . Tissue Specificity Lung, Function function:This protein aggregates to form microtubular structures.,induction:By alpha and beta interferons, but not by gamma interferons.,similarity:Belongs to the IFI44 family., Background function:This protein aggregates to form microtubular structures.,induction:By alpha and beta interferons, but not by gamma interferons.,similarity:Belongs to the IFI44 family., matters needing Avoid repeated freezing and thawing!	Storage Stability	-20°C/1 year
Cell Pathway Cytoplasm . Lung, function function:This protein aggregates to form microtubular structures.,induction:By alpha and beta interferons, but not by gamma interferons.,similarity:Belongs to the IFI44 family., Background function:This protein aggregates to form microtubular structures.,induction:By alpha and beta interferons, but not by gamma interferons.,similarity:Belongs to the IFI44 family., matters needing Avoid repeated freezing and thawing!	Synonyms	
Tissue Specificity Lung, function function: This protein aggregates to form microtubular structures., induction: By alpha and beta interferons, but not by gamma interferons., similarity: Belongs to the IFI44 family., Background function: This protein aggregates to form microtubular structures., induction: By alpha and beta interferons, but not by gamma interferons., similarity: Belongs to the IFI44 family., matters needing Avoid repeated freezing and thawing!	Observed Band	48kD
Function function: This protein aggregates to form microtubular structures.,induction: By alpha and beta interferons, but not by gamma interferons.,similarity: Belongs to the IFI44 family., function: This protein aggregates to form microtubular structures.,induction: By alpha and beta interferons, but not by gamma interferons.,similarity: Belongs to the IFI44 family., matters needing Avoid repeated freezing and thawing!	Cell Pathway	Cytoplasm .
alpha and beta interferons, but not by gamma interferons.,similarity:Belongs to the IFI44 family., Background function:This protein aggregates to form microtubular structures.,induction:By alpha and beta interferons, but not by gamma interferons.,similarity:Belongs to the IFI44 family., matters needing Avoid repeated freezing and thawing!	Tissue Specificity	Lung,
alpha and beta interferons, but not by gamma interferons.,similarity:Belongs to the IFI44 family., matters needing Avoid repeated freezing and thawing!	Function	alpha and beta interferons, but not by gamma interferons, similarity: Belongs to the
matters needing Avoid repeated freezing and thawing! attention	Background	alpha and beta interferons, but not by gamma interferons, similarity: Belongs to the
	matters needing attention	Avoid repeated freezing and thawing!

Nanjing BYabscience technology Co.,Ltd



国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询



ı	Isaq	P S	ua	ae	stic	าทร
u	Jau	C 3	uч	uc.	วแง	ノロコ

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images							

Nanjing BYabscience technology Co.,Ltd