



KDM5A Polyclonal Antibody

Catalog No	BYab-05113
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB;ELISA
Gene Name	KDM5A JARID1A RBBP2 RBP2
Protein Name	Lysine-specific demethylase 5A (EC 1.14.11.-) (Histone demethylase JARID1A) (Jumonji/ARID domain-containing protein 1A) (Retinoblastoma-binding protein 2) (RBBP-2)
Immunogen	Synthesized peptide derived from human protein . at AA range: 720-800
Specificity	KDM5A Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	185kD
Cell Pathway	Nucleus, nucleolus . Nucleus . Occupies promoters of genes involved in RNA metabolism and mitochondrial function. .
Tissue Specificity	Epithelium, Eye, Testis,
Function	cofactor:Fe(2+).,function:Histone demethylase that specifically demethylates 'Lys-4' of histone H3, thereby playing a central role in histone code. Does not demethylate histone H3 'Lys-9', H3 'Lys-27', H3 'Lys-36', H3 'Lys-79' or H4 'Lys-20'. Demethylates trimethylated and dimethylated but not monomethylated H3 'Lys-4'. May stimulate transcription mediated by nuclear receptors. May be involved in transcriptional regulation of Hox proteins during cell differentiation. May participate in transcriptional repression of cytokines such as CXCL12.,similarity:Belongs to the JARID1 histone demethylase family.,similarity:Contains 1 ARID domain.,similarity:Contains 1 JmjC domain.,similarity:Contains 1 JmjN domain.,similarity:Contains 3 PHD-type zinc

Nanjing BYabscience technology Co.,Ltd



fingers.,subunit:Interacts with the viral protein-binding domain of RB1. Interacts with ESR1, MYC, MYCN and LMO2.,

Background

This gene encodes a member of the Jumonji, AT-rich interactive domain 1 (JARID1) histone demethylase protein family. The encoded protein plays a role in gene regulation through the histone code by specifically demethylating lysine 4 of histone H3. The encoded protein interacts with many other proteins, including retinoblastoma protein, and is implicated in the transcriptional regulation of Hox genes and cytokines. This gene may play a role in tumor progression. [provided by RefSeq, Aug 2013],

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images