



MX1 Polyclonal Antibody

Catalog No	BYab-05138
Isotype	IgG
Reactivity	Human;Rat
Applications	WB;ELISA
Gene Name	MX1
Protein Name	Interferon-induced GTP-binding protein Mx1 (Interferon-induced protein p78) (IFI-78K) (Interferon-regulated resistance GTP-binding protein MxA) (Myxoma resistance protein 1) (Myxovirus resistance prot
Immunogen	Synthesized peptide derived from human protein . at AA range: 500-580
Specificity	MX1 Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	72kD
Cell Pathway	Cytoplasm . Endoplasmic reticulum membrane ; Peripheral membrane protein ; Cytoplasmic side. Cytoplasm, perinuclear region . Binds preferentially to negatively charged phospholipids (PubMed:21900240). Colocalizes with CCHFV protein N in the perinuclear region (PubMed:15047845). . ; [Isoform 2]: Cytoplasm . Nucleus . Translocates into the nuclei of HSV-1 infected cells (PubMed:20603636). .
Tissue Specificity	Primary B-Cells,Spleen,Uterus,
Function	function:May regulate the calcium channel activity of TRPCs. Shows activity against influenza virus and VSV, a rhabdovirus.,induction:By interferons.,similarity:Belongs to the dynamin family.,similarity:Contains 1 GED domain.,subunit:Interacts with TRPC1, TRPC3, TRPC4, TRPC5, TRPC6 and TRPC7.,

Nanjing BYabscience technology Co.,Ltd



Background

This gene encodes a guanosine triphosphate (GTP)-metabolizing protein that participates in the cellular antiviral response. The encoded protein is induced by type I and type II interferons and antagonizes the replication process of several different RNA and DNA viruses. There is a related gene located adjacent to this gene on chromosome 21, and there are multiple pseudogenes located in a cluster on chromosome 4. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Sep 2013],

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images