



GOT1B Polyclonal Antibody

Catalog No	BYab-05194
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB;ELISA
Gene Name	GOLT1B GOT1A CGI-141 HDCMA39P UNQ432/PRO793
Protein Name	Vesicle transport protein GOT1B (Golgi transport 1 homolog B) (Putative NF-kappa-B-activating protein 470) (hGOT1a)
Immunogen	Synthesized peptide derived from human protein . at AA range: 70-150
Specificity	GOT1B Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Concentration	: 111 9 /1111
Purity	≥90%
	•
Purity	≥90%
Purity Storage Stability	≥90%
Purity Storage Stability Synonyms	≥90% -20°C/1 year
Purity Storage Stability Synonyms Observed Band	≥90% -20°C/1 year 15kD
Purity Storage Stability Synonyms Observed Band Cell Pathway	≥90% -20°C/1 year 15kD Golgi apparatus membrane ; Multi-pass membrane protein . Widely expressed. Tends to be up-regulated in seminomas compared to normal
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity	≥90% -20°C/1 year 15kD Golgi apparatus membrane ; Multi-pass membrane protein . Widely expressed. Tends to be up-regulated in seminomas compared to normal testis. function:May be involved in fusion of ER-derived transport vesicles with the Golgi
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function	≥90% -20°C/1 year 15kD Golgi apparatus membrane ; Multi-pass membrane protein . Widely expressed. Tends to be up-regulated in seminomas compared to normal testis. function:May be involved in fusion of ER-derived transport vesicles with the Golgi complex.,similarity:Belongs to the GOT1 family., function:May be involved in fusion of ER-derived transport vesicles with the Golgi
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function Background matters needing	≥90% -20°C/1 year 15kD Golgi apparatus membrane; Multi-pass membrane protein. Widely expressed. Tends to be up-regulated in seminomas compared to normal testis. function:May be involved in fusion of ER-derived transport vesicles with the Golgi complex.,similarity:Belongs to the GOT1 family., function:May be involved in fusion of ER-derived transport vesicles with the Golgi complex.,similarity:Belongs to the GOT1 family.,


Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658





Products Images



Western blot analysis of lysates from U2OS cells, primary antibody was diluted at 1:1000, 4°over night

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658