




KLF16 Polyclonal Antibody

Catalog No BYab-05201 Isotype IgG Reactivity Human;Mouse Applications WB;ELISA Gene Name KLF16 BTEB4 NSLP2 Protein Name Krueppel-like factor 16 (Basic transcription element-binding protein 4) (Novel Sp1-like zinc finger transcription factor 2) (Transcription factor BTEB4) (Transcription factor N Immunogen Synthesized peptide derived from human protein . at AA range: 10-90 Specificity KLF16 Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit.lgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 27kD Cell Pathway Nucleus Tissue Specificity Pancreas, Function function:Transcription factor that binds GC and GT boxes and displaces Sp1 and Sp3 from these sequences. Modulates dopaminergic transmission in the brain		
Reactivity Human;Mouse Applications WB;ELISA Gene Name KLF16 BTEB4 NSLP2 Protein Name Krueppel-like factor 16 (Basic transcription element-binding protein 4) (Novel Sp1-like zinc finger transcription factor 2) (Transcription factor BTEB4) (Transcription factor N Immunogen Synthesized peptide derived from human protein . at AA range: 10-90 Specificity KLF16 Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 27kD Cell Pathway Nucleus . Tissue Specificity Pancreas, Function function:Transcription factor that binds GC and GT boxes and displaces Sp1 and Sp3 from these sequences. Modulates dopaminergic transmission in the brain, similarity:Belongs to the Sp1 C2H2-type zinc-finger protein family, similarity:Belongs to the Sp1 C2H2-type zinc-finger protein	Catalog No	BYab-05201
Applications WB;ELISA Gene Name KLF16 BTEB4 NSLP2 Protein Name Krueppel-like factor 16 (Basic transcription element-binding protein 4) (BTE-binding protein 4) (Novel Sp1-like zinc finger transcription factor 2) (Transcription factor BTEB4) (Transcription factor N Immunogen Synthesized peptide derived from human protein . at AA range: 10-90 Specificity KLF16 Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 27kD Cell Pathway Nucleus . Tissue Specificity Pancreas, Function function:Transcription factor that binds GC and GT boxes and displaces Sp1 and Sp3 from these sequences. Modulates dopaminergic transmission in the brain similarity. Belongs to the Sp1 C2PI2-type zinc-finger protein family, similarity. Contains 3 C2H2-type zinc-finger protein brain is protein for the brain similarity. Belongs to the Sp1 C2PI2-type zinc-finger protein family brain. Sp3 from these sequences. Modulates dopaminergic transmission in the brain similarity. Belongs to the Sp1 C2PI2-type zinc-finger protein family brain. Sp3 from these sequences. Modulates dopaminergic transmission in the brain similarity. Belongs to the Sp1 C2PI2-type zinc-finger protein family brain. Sp1 C2PI2-type zinc-finger protein family brain. Sp1 C2PI2-type zinc-finger protein family spain.	Isotype	IgG
Gene Name KLF16 BTEB4 NSLP2 Protein Name Krueppel-like factor 16 (Basic transcription element-binding protein 4) (Novel Sp1-like zinc finger transcription factor 2) (Transcription factor BTEB4) (Transcription factor N Immunogen Synthesized peptide derived from human protein . at AA range: 10-90 Specificity KLF16 Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, lgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 27kD Cell Pathway Nucleus . Tissue Specificity Pancreas, Function function:Transcription factor that binds GC and GT boxes and displaces Sp1 and Sp3 from these sequences. Modulates dopaminergic transmission in the brain, similarity:Belongs to the Sp1 C2H2-type zinc-finger protein family, similarity:Contains 3 C2H2-type zinc-finger protein in the brain, similarity:Belongs to the Sp1 C2H2-type zinc-finger protein	Reactivity	Human;Mouse
Protein Name Krueppel-like factor 16 (Basic transcription element-binding protein 4) (BTE-binding protein 4) (Novel Sp1-like zinc finger transcription factor 2) (Transcription factor BTEB4) (Transcription factor N Immunogen Synthesized peptide derived from human protein . at AA range: 10-90 Specificity KLF16 Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 27kD Cell Pathway Nucleus Tissue Specificity Pancreas, Function function:Transcription factor that binds GC and GT boxes and displaces Sp1 and Sp3 from these sequences. Modulates dopaminergic transmission in the brain, similarity:Belongs to the Sp1 C2H2-type zinc-finger protein family, similarity:Gongton factor that binds GC and GT boxes and displaces Sp1 and Sp3 from these sequences. Modulates dopaminergic transmission in the brain, similarity:Belongs to the Sp1 C2H2-type zinc-finger protein	Applications	WB;ELISA
Immunogen Synthesized peptide derived from human protein . at AA range: 10-90 Specificity KLF16 Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 27kD Cell Pathway Nucleus . Tissue Specificity Pancreas, Function function:Transcription factor that binds GC and GT boxes and displaces Sp1 and Sp3 from these sequences. Modulates dopaminergic transmission in the brain.,similarity.Belongs to the Sp1 C2H2-type zinc-finger protein family, similarity.Contains 3 C2H2-type zinc-fingers. Background function:Transcription factor that binds GC and GT boxes and displaces Sp1 and Sp3 from these sequences. Modulates dopaminergic transmission in the brain.,similarity.Belongs to the Sp1 C2H2-type zinc-finger protein in the brain.,similarity.Belongs to the Sp1 C2H2-type zinc-finger protein	Gene Name	KLF16 BTEB4 NSLP2
Specificity KLF16 Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Nucleus . Tissue Specificity Pancreas, Function function:Transcription factor that binds GC and GT boxes and displaces Sp1 and Sp3 from these sequences. Modulates dopaminergic transmission in the brain.,similarity.Contains 3 C2H2-type zinc finger protein family ,similarity.Contains 3 C2H2-type zinc finger protein family ,similarity.Contains 3 C2H2-type zinc finger protein family ,similarity.Eelongs to the Sp1 C2H2-type zinc-finger protein family ,similarity.Eelongs to the Sp1 C2H2-type zinc-finger protein	Protein Name	Krueppel-like factor 16 (Basic transcription element-binding protein 4) (BTE-binding protein 4) (Novel Sp1-like zinc finger transcription factor 2) (Transcription factor BTEB4) (Transcription factor N
Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Nucleus . Tissue Specificity Pancreas, Function function:Transcription factor that binds GC and GT boxes and displaces Sp1 and Sp3 from these sequences. Modulates dopaminergic transmission in the brain. similarity:Belongs to the Sp1 C2H2-type zinc-finger protein familysimilarity:Contains 3 C2H2-type zinc fingers., Background function:Transcription factor that binds GC and GT boxes and displaces Sp1 and Sp3 from these sequences. Modulates dopaminergic transmission in the brainsimilarity:Belongs to the Sp1 C2H2-type zinc-finger protein	Immunogen	Synthesized peptide derived from human protein . at AA range: 10-90
Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 27kD Cell Pathway Nucleus . Tissue Specificity Pancreas, Function function:Transcription factor that binds GC and GT boxes and displaces Sp1 and Sp3 from these sequences. Modulates dopaminergic transmission in the brain., similarity:Belongs to the Sp1 C2H2-type zinc-finger protein family., similarity:Contains 3 C2H2-type zinc fingers., Background function:Transcription factor that binds GC and GT boxes and displaces Sp1 and Sp3 from these sequences. Modulates dopaminergic transmission in the brain., similarity:Belongs to the Sp1 C2H2-type zinc-finger protein	Specificity	KLF16 Polyclonal Antibody detects endogenous levels of protein.
Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 27kD Cell Pathway Nucleus . Tissue Specificity Pancreas, Function function:Transcription factor that binds GC and GT boxes and displaces Sp1 and Sp3 from these sequences. Modulates dopaminergic transmission in the brain.,similarity:Belongs to the Sp1 C2H2-type zinc-finger protein family, similarity:Contains 3 C2H2-type zinc fingers., Background function:Transcription factor that binds GC and GT boxes and displaces Sp1 and Sp3 from these sequences. Modulates dopaminergic transmission in the brain, similarity:Belongs to the Sp1 C2H2-type zinc-finger protein in the brain, similarity:Belongs to the Sp1 C2H2-type zinc-finger protein	Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 27kD Cell Pathway Nucleus . Tissue Specificity Pancreas, Function function:Transcription factor that binds GC and GT boxes and displaces Sp1 and Sp3 from these sequences. Modulates dopaminergic transmission in the brain.,similarity:Belongs to the Sp1 C2H2-type zinc-finger protein family.,similarity:Contains 3 C2H2-type zinc fingers., Background function:Transcription factor that binds GC and GT boxes and displaces Sp1 and Sp3 from these sequences. Modulates dopaminergic transmission in the brain.,similarity:Belongs to the Sp1 C2H2-type zinc-finger protein family.,similarity:Belongs to the Sp1 C2H2-type zinc-finger protein	Source	Polyclonal, Rabbit,IgG
Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms -27kD Cell Pathway Nucleus . Tissue Specificity Pancreas, Function function:Transcription factor that binds GC and GT boxes and displaces Sp1 and Sp3 from these sequences. Modulates dopaminergic transmission in the brain., similarity:Belongs to the Sp1 C2H2-type zinc-finger protein family., similarity:Contains 3 C2H2-type zinc fingers., Background function:Transcription factor that binds GC and GT boxes and displaces Sp1 and Sp3 from these sequences. Modulates dopaminergic transmission in the brain., similarity:Belongs to the Sp1 C2H2-type zinc-finger protein	Purification	·
Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 27kD Cell Pathway Nucleus . Tissue Specificity Pancreas, Function function:Transcription factor that binds GC and GT boxes and displaces Sp1 and Sp3 from these sequences. Modulates dopaminergic transmission in the brain.,similarity:Belongs to the Sp1 C2H2-type zinc-finger protein family.,similarity:Contains 3 C2H2-type zinc fingers., Background function:Transcription factor that binds GC and GT boxes and displaces Sp1 and Sp3 from these sequences. Modulates dopaminergic transmission in the brain.,similarity:Belongs to the Sp1 C2H2-type zinc fingers.,	Dilution	WB 1:500-2000 ELISA 1:5000-20000
Storage Stability -20°C/1 year Synonyms Observed Band 27kD Cell Pathway Nucleus . Tissue Specificity Pancreas, Function function:Transcription factor that binds GC and GT boxes and displaces Sp1 and Sp3 from these sequences. Modulates dopaminergic transmission in the brain.,similarity:Belongs to the Sp1 C2H2-type zinc-finger protein family.,similarity:Contains 3 C2H2-type zinc fingers., Background function:Transcription factor that binds GC and GT boxes and displaces Sp1 and Sp3 from these sequences. Modulates dopaminergic transmission in the brain.,similarity:Belongs to the Sp1 C2H2-type zinc-finger protein	Concentration	1 mg/ml
Synonyms Observed Band 27kD Cell Pathway Nucleus . Tissue Specificity Pancreas, Function function:Transcription factor that binds GC and GT boxes and displaces Sp1 and Sp3 from these sequences. Modulates dopaminergic transmission in the brain.,similarity:Belongs to the Sp1 C2H2-type zinc-finger protein family.,similarity:Contains 3 C2H2-type zinc fingers., Background function:Transcription factor that binds GC and GT boxes and displaces Sp1 and Sp3 from these sequences. Modulates dopaminergic transmission in the brain.,similarity:Belongs to the Sp1 C2H2-type zinc-finger protein	Purity	≥90%
Observed Band 27kD Cell Pathway Nucleus . Tissue Specificity Pancreas, Function function:Transcription factor that binds GC and GT boxes and displaces Sp1 and Sp3 from these sequences. Modulates dopaminergic transmission in the brain.,similarity:Belongs to the Sp1 C2H2-type zinc-finger protein family.,similarity:Contains 3 C2H2-type zinc fingers., Background function:Transcription factor that binds GC and GT boxes and displaces Sp1 and Sp3 from these sequences. Modulates dopaminergic transmission in the brain.,similarity:Belongs to the Sp1 C2H2-type zinc-finger protein	Storage Stability	-20°C/1 year
Cell Pathway Nucleus . Tissue Specificity Pancreas, Function function:Transcription factor that binds GC and GT boxes and displaces Sp1 and Sp3 from these sequences. Modulates dopaminergic transmission in the brain.,similarity:Belongs to the Sp1 C2H2-type zinc-finger protein family.,similarity:Contains 3 C2H2-type zinc fingers., Background function:Transcription factor that binds GC and GT boxes and displaces Sp1 and Sp3 from these sequences. Modulates dopaminergic transmission in the brain.,similarity:Belongs to the Sp1 C2H2-type zinc-finger protein	Synonyms	
Tissue Specificity Pancreas, Function function:Transcription factor that binds GC and GT boxes and displaces Sp1 and Sp3 from these sequences. Modulates dopaminergic transmission in the brain.,similarity:Belongs to the Sp1 C2H2-type zinc-finger protein family.,similarity:Contains 3 C2H2-type zinc fingers., Background function:Transcription factor that binds GC and GT boxes and displaces Sp1 and Sp3 from these sequences. Modulates dopaminergic transmission in the brain.,similarity:Belongs to the Sp1 C2H2-type zinc-finger protein	Observed Band	27kD
Function function:Transcription factor that binds GC and GT boxes and displaces Sp1 and Sp3 from these sequences. Modulates dopaminergic transmission in the brain.,similarity:Belongs to the Sp1 C2H2-type zinc-finger protein family.,similarity:Contains 3 C2H2-type zinc fingers., function:Transcription factor that binds GC and GT boxes and displaces Sp1 and Sp3 from these sequences. Modulates dopaminergic transmission in the brain.,similarity:Belongs to the Sp1 C2H2-type zinc-finger protein	Cell Pathway	Nucleus .
Sp3 from these sequences. Modulates dopaminergic transmission in the brain.,similarity:Belongs to the Sp1 C2H2-type zinc-finger protein family.,similarity:Contains 3 C2H2-type zinc fingers., Background function:Transcription factor that binds GC and GT boxes and displaces Sp1 and Sp3 from these sequences. Modulates dopaminergic transmission in the brain.,similarity:Belongs to the Sp1 C2H2-type zinc-finger protein	Tissue Specificity	Pancreas,
Sp3 from these sequences. Modulates dopaminergic transmission in the brain.,similarity:Belongs to the Sp1 C2H2-type zinc-finger protein	Function	Sp3 from these sequences. Modulates dopaminergic transmission in the brain similarity:Belongs to the Sp1 C2H2-type zinc-finger protein
	Background	Sp3 from these sequences. Modulates dopaminergic transmission in the brain.,similarity:Belongs to the Sp1 C2H2-type zinc-finger protein

Nanjing BYabscience technology Co.,Ltd








matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Nanjing BYabscience technology Co.,Ltd