



MOGT3 Polyclonal Antibody

Catalog No	BYab-05222
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB;ELISA
Gene Name	MOGAT3 DC7 DGAT2L7 UNQ9383/PRO34208
Protein Name	2-acylglycerol O-acyltransferase 3 (EC 2.3.1.20) (EC 2.3.1.22) (Acyl-CoA:monoacylglycerol acyltransferase 3) (MGAT3) (Diacylglycerol O-acyltransferase candidate 7) (hDC7) (Diacylglycerol acyltransferase)
Immunogen	Synthesized peptide derived from human protein . at AA range: 10-90
Specificity	MOGT3 Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	37kD
Cell Pathway	Endoplasmic reticulum membrane ; Multi-pass membrane protein . Cytoplasm, perinuclear region .
Tissue Specificity	Selectively expressed in the digestive system. Highly expressed in the ileum, and at lower level in jejunum, duodenum, colon, cecum and the rectum. Not expressed in the stomach and the esophagus and trachea. Expressed at very low level in liver.
Function	catalytic activity:Acyl-CoA + 1,2-diacylglycerol = CoA + triacylglycerol.,catalytic activity:Acyl-CoA + 2-acylglycerol = CoA + diacylglycerol.,function:Catalyzes the formation of diacylglycerol from 2-monoacylglycerol and fatty acyl-CoA. Also able to catalyze the terminal step in triacylglycerol synthesis by using diacylglycerol and fatty acyl CoA as substrates. Has a preference toward palmitoyl-CoA and oleoyl-CoA. May be involved in absorption of dietary fat in the small intestine by catalyzing the resynthesis of triacylglycerol in enterocytes.,pathway:Glycerolipid metabolism; triacylglycerol biosynthesis.,similarity:Belongs to the diacylglycerol

Nanjing BYabscience technology Co.,Ltd



acyltransferase family.,tissue specificity:Selectively expressed in the digestive system. Highly expressed in the ileum, and at lower level in jejunum, duodenum, colon, cecum and the rectum. Not expressed in the stomach and the esophagus and tr

Background

Acyl-CoA:monoacylglycerol acyltransferase (MOGAT; EC 2.3.1.22) catalyzes the synthesis of diacylglycerol from 2-monoacylglycerol and fatty acyl-CoA (Cheng et al., 2003 [PubMed 12618427]).[supplied by OMIM, Mar 2008],

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

