



## ABHDA Polyclonal Antibody

hydrolase. The encoded protein removes glucuronide from mycophenolic acid acyl-glucuronide. There is a pseudogene for this gene on chromosome 6.		
Reactivity Human;Rat  Applications WB:ELISA  Gene Name ABHD10  Protein Name Abhydrolase domain-containing protein 10, mitochondrial (EC 3.4)  Immunogen Synthesized peptide derived from human protein . at AA range: 220-300  Specificity ABHDA Polyclonal Antibody detects endogenous levels of protein.  Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.  Source Polyclonal, Rabbit,IgG  Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000 ELISA 1:5000-20000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band 33kD  Cell Pathway Mitochondrion.  Tissue Specificity Melanoma,Ovary,Placenta,  Function similarity:Belongs to the AB hydrolase superfamily.,  Background This gene encodes a mitochondrially-localized enzyme that acts in liver cells as a hydrolase. The encoded protein removes glucuronide from mycophenolic acid acyl-glucuronide. There is a pseudogene for this gene on chromosome 6, Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jan 2013],  matters needing Avoid repeated freezing and thawing!	Catalog No	BYab-05262
Applications WB;ELISA  Gene Name ABHD10  Protein Name Abhydrolase domain-containing protein 10, mitochondrial (EC 3.4)  Immunogen Synthesized peptide derived from human protein . at AA range: 220-300  Specificity ABHDA Polyclonal Antibody detects endogenous levels of protein.  Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.  Source Polyclonal, Rabbit,IgG  Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000 ELISA 1:5000-20000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band 33kD  Cell Pathway Mitochondrion .  Tissue Specificity Melanoma,Ovary,Placenta,  Function similarity:Belongs to the AB hydrolase superfamily.,  Background This gene encodes a mitochondrially-localized enzyme that acts in liver cells as a hydrolase. The encoded protein removes glucuronide from mycophenolic acid acty-glucuronide. There is a pseudogene for this gene on chromosome 6. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jan 2013],  matters needing Avoid repeated freezing and thawing!	Isotype	IgG
Gene Name ABHD10 Protein Name Abhydrolase domain-containing protein 10, mitochondrial (EC 3.4) Immunogen Synthesized peptide derived from human protein . at AA range: 220-300 Specificity ABHDA Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 33kD Cell Pathway Mitochondrion . Tissue Specificity Melanoma,Ovary,Placenta, Function similarity:Belongs to the AB hydrolase superfamily., Background This gene encodes a mitochondrially-localized enzyme that acts in liver cells as a hydrolase. The encoded protein removes glucuronide from mycophenolic acid acty-glucuronide. There is a pseudogene for this gene on chromosome 6, Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jan 2013], matters needing Avoid repeated freezing and thawing!	Reactivity	Human;Rat
Protein Name Abhydrolase domain-containing protein 10, mitochondrial (EC 3.4)  Immunogen Synthesized peptide derived from human protein . at AA range: 220-300  Specificity ABHDA Polyclonal Antibody detects endogenous levels of protein.  Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.  Source Polyclonal, Rabbit,IgG  Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000 ELISA 1:5000-20000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band 33kD  Cell Pathway Mitochondrion .  Tissue Specificity Melanoma,Ovary,Placenta,  Function similarity:Belongs to the AB hydrolase superfamily.,  This gene encodes a mitochondrially-localized enzyme that acts in liver cells as a hydrolase. The encoded protein removes glucuronide from mycophenolic acid acid-acyl-glucuronide. There is a pseudogene for this gene on chromosome 6. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jan 2013],  matters needing Avoid repeated freezing and thawing!	Applications	WB;ELISA
Immunogen       Synthesized peptide derived from human protein . at AA range: 220-300         Specificity       ABHDA Polyclonal Antibody detects endogenous levels of protein.         Formulation       Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.         Source       Polyclonal, Rabbit, IgG         Purification       The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.         Dilution       WB 1:500-2000 ELISA 1:5000-20000         Concentration       1 mg/ml         Purity       ≥90%         Storage Stability       -20°C/1 year         Synonyms       Observed Band         Observed Band       33kD         Cell Pathway       Mitochondrion .         Tissue Specificity       Melanoma,Ovary,Placenta,         Function       similarity:Belongs to the AB hydrolase superfamily.,         Background       This gene encodes a mitochondrially-localized enzyme that acts in liver cells as a hydrolase. The encoded protein removes glucuronide from mycophenolic acid acyt-glucuronide. There is a pseudogene for this gene on chromosome 6. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jan 2013],         matters needing       Avoid repeated freezing and thawing!	Gene Name	ABHD10
Specificity  ABHDA Polyclonal Antibody detects endogenous levels of protein.  Formulation  Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.  Source  Polyclonal, Rabbit, IgG  Purification  The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution  WB 1:500-2000 ELISA 1:5000-20000  Concentration  1 mg/ml  Purity  ≥90%  Storage Stability  -20°C/1 year  Synonyms  Observed Band  33kD  Cell Pathway  Mitochondrion .  Tissue Specificity  Melanoma, Ovary, Placenta,  Function  Similarity: Belongs to the AB hydrolase superfamily.,  Background  This gene encodes a mitochondrially-localized enzyme that acts in liver cells as a hydrolase. The encoded protein removes glucuronide from mycophenolic acid acyl-glucuronide. There is a pseudogene for this gene on chromosome 6, Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jan 2013],  matters needing  Avoid repeated freezing and thawing!	Protein Name	Abhydrolase domain-containing protein 10, mitochondrial (EC 3.4)
Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.  Source Polyclonal, Rabbit,IgG  Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000 ELISA 1:5000-20000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band 33kD  Cell Pathway Mitochondrion .  Tissue Specificity Melanoma,Ovary,Placenta,  Function similarity:Belongs to the AB hydrolase superfamily.,  This gene encodes a mitochondrially-localized enzyme that acts in liver cells as a hydrolase. The encoded protein removes glucuronide from mycophenolic acid acyl-glucuronide. There is a pseudogene for this gene on chromosome 6. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jan 2013],  matters needing Avoid repeated freezing and thawing!	Immunogen	Synthesized peptide derived from human protein . at AA range: 220-300
Source       Polyclonal, Rabbit,IgG         Purification       The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.         Dilution       WB 1:500-2000 ELISA 1:5000-20000         Concentration       1 mg/ml         Purity       ≥90%         Storage Stability       -20°C/1 year         Synonyms       Observed Band       33kD         Cell Pathway       Mitochondrion .         Tissue Specificity       Melanoma,Ovary,Placenta,         Function       similarity:Belongs to the AB hydrolase superfamily.,         Background       This gene encodes a mitochondrially-localized enzyme that acts in liver cells as a hydrolase. The encoded protein removes glucuronide from mycophenolic acid acyl-glucuronide. There is a pseudogene for this gene on chromosome 6. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jan 2013],         matters needing       Avoid repeated freezing and thawing!	Specificity	ABHDA Polyclonal Antibody detects endogenous levels of protein.
Purification  The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution  WB 1:500-2000 ELISA 1:5000-20000  Concentration  1 mg/ml  Purity  ≥90%  Storage Stability  -20°C/1 year  Synonyms  Observed Band  33kD  Cell Pathway  Mitochondrion .  Tissue Specificity  Melanoma,Ovary,Placenta,  Function  similarity:Belongs to the AB hydrolase superfamily.,  This gene encodes a mitochondrially-localized enzyme that acts in liver cells as a hydrolase. The encoded protein removes glucuronide from mycophenolic acid acyl-glucuronide. There is a pseudogene for this gene on chromosome 6. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jan 2013],  matters needing  Avoid repeated freezing and thawing!	Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000 ELISA 1:5000-20000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band 33kD  Cell Pathway Mitochondrion .  Tissue Specificity Melanoma,Ovary,Placenta,  Function similarity:Belongs to the AB hydrolase superfamily.,  Background This gene encodes a mitochondrially-localized enzyme that acts in liver cells as a hydrolase. The encoded protein removes glucuronide from mycophenolic acid acyl-glucuronide. There is a pseudogene for this gene on chromosome 6. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jan 2013],  matters needing Avoid repeated freezing and thawing!	Source	Polyclonal, Rabbit,IgG
Concentration       1 mg/ml         Purity       ≥90%         Storage Stability       -20°C/1 year         Synonyms       -20°C/1 year         Observed Band       33kD         Cell Pathway       Mitochondrion .         Tissue Specificity       Melanoma,Ovary,Placenta,         Function       similarity:Belongs to the AB hydrolase superfamily.,         Background       This gene encodes a mitochondrially-localized enzyme that acts in liver cells as a hydrolase. The encoded protein removes glucuronide from mycophenolic acid acyl-glucuronide. There is a pseudogene for this gene on chromosome 6. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jan 2013],         matters needing       Avoid repeated freezing and thawing!	Purification	
Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band 33kD  Cell Pathway Mitochondrion .  Tissue Specificity Melanoma,Ovary,Placenta,  Function similarity:Belongs to the AB hydrolase superfamily.,  Background This gene encodes a mitochondrially-localized enzyme that acts in liver cells as a hydrolase. The encoded protein removes glucuronide from mycophenolic acid acyl-glucuronide. There is a pseudogene for this gene on chromosome 6. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jan 2013],  matters needing Avoid repeated freezing and thawing!	Dilution	WB 1:500-2000 ELISA 1:5000-20000
Storage Stability -20°C/1 year  Synonyms  Observed Band 33kD  Cell Pathway Mitochondrion .  Tissue Specificity Melanoma,Ovary,Placenta,  Function similarity:Belongs to the AB hydrolase superfamily.,  Background This gene encodes a mitochondrially-localized enzyme that acts in liver cells as a hydrolase. The encoded protein removes glucuronide from mycophenolic acid acyl-glucuronide. There is a pseudogene for this gene on chromosome 6. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jan 2013],  matters needing Avoid repeated freezing and thawing!	Concentration	1 mg/ml
Synonyms  Observed Band 33kD  Cell Pathway Mitochondrion .  Tissue Specificity Melanoma,Ovary,Placenta,  Function similarity:Belongs to the AB hydrolase superfamily.,  Background This gene encodes a mitochondrially-localized enzyme that acts in liver cells as a hydrolase. The encoded protein removes glucuronide from mycophenolic acid acyl-glucuronide. There is a pseudogene for this gene on chromosome 6. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jan 2013],  matters needing Avoid repeated freezing and thawing!	Purity	≥90%
Observed Band  Cell Pathway  Mitochondrion .  Tissue Specificity  Melanoma,Ovary,Placenta,  Function  Similarity:Belongs to the AB hydrolase superfamily.,  Background  This gene encodes a mitochondrially-localized enzyme that acts in liver cells as a hydrolase. The encoded protein removes glucuronide from mycophenolic acid acyl-glucuronide. There is a pseudogene for this gene on chromosome 6. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jan 2013],  matters needing  Avoid repeated freezing and thawing!	Storage Stability	-20°C/1 year
Cell Pathway  Mitochondrion .  Tissue Specificity  Melanoma, Ovary, Placenta,  Function  Similarity: Belongs to the AB hydrolase superfamily.,  This gene encodes a mitochondrially-localized enzyme that acts in liver cells as a hydrolase. The encoded protein removes glucuronide from mycophenolic acid acyl-glucuronide. There is a pseudogene for this gene on chromosome 6. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jan 2013],  matters needing  Avoid repeated freezing and thawing!	Synonyms	
Tissue Specificity  Melanoma,Ovary,Placenta,  Function  similarity:Belongs to the AB hydrolase superfamily.,  This gene encodes a mitochondrially-localized enzyme that acts in liver cells as a hydrolase. The encoded protein removes glucuronide from mycophenolic acid acyl-glucuronide. There is a pseudogene for this gene on chromosome 6. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jan 2013],  matters needing  Avoid repeated freezing and thawing!	Observed Band	33kD
Function similarity:Belongs to the AB hydrolase superfamily.,  Background This gene encodes a mitochondrially-localized enzyme that acts in liver cells as a hydrolase. The encoded protein removes glucuronide from mycophenolic acid acyl-glucuronide. There is a pseudogene for this gene on chromosome 6. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jan 2013],  matters needing Avoid repeated freezing and thawing!	Cell Pathway	Mitochondrion .
Background  This gene encodes a mitochondrially-localized enzyme that acts in liver cells as a hydrolase. The encoded protein removes glucuronide from mycophenolic acid acyl-glucuronide. There is a pseudogene for this gene on chromosome 6. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jan 2013],  matters needing  Avoid repeated freezing and thawing!	Tissue Specificity	Melanoma,Ovary,Placenta,
hydrolase. The encoded protein removes glucuronide from mycophenolic acid acyl-glucuronide. There is a pseudogene for this gene on chromosome 6. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jan 2013],  matters needing  Avoid repeated freezing and thawing!	Function	similarity:Belongs to the AB hydrolase superfamily.,
matters needing Avoid repeated freezing and thawing! attention	Background	acyl-glucuronide. There is a pseudogene for this gene on chromosome 6.  Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jan
	matters needing attention	Avoid repeated freezing and thawing!

Nanjing BYabscience technology Co.,Ltd





**Usage suggestions** 

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

