




## ATX10 Polyclonal Antibody

| Catalog NoBYab-05337IsotypeIgGReactivityHuman;Mouse;RatApplicationsWB;ELISAGene NameATXN10 SCA10Protein NameAtaxin-10 (Brain protein E46 homolog) (Spinocerebellar ataxia type 10ImmunogenSynthesized peptide derived from human protein . at AA range: 10-90SpecificityATX10 Polyclonal Antibody detects endogenous levels of protein.FormulationLiquid in PBS containing 50% glycerol, and 0.02% sodium azide.SourcePolyclonal, Rabbit,IgGPurificationThe antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.DilutionWB 1:500-2000 ELISA 1:5000-20000Concentration1 mg/ml | protein)                         |
|---|----------------------------------|
| Reactivity Human;Mouse;Rat  Applications WB;ELISA  Gene Name ATXN10 SCA10  Protein Name Ataxin-10 (Brain protein E46 homolog) (Spinocerebellar ataxia type 10 lmmunogen Synthesized peptide derived from human protein . at AA range: 10-90  Specificity ATX10 Polyclonal Antibody detects endogenous levels of protein.  Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.  Source Polyclonal, Rabbit,IgG  Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000 ELISA 1:5000-20000                        | protein)                         |
| Applications WB;ELISA  Gene Name ATXN10 SCA10  Protein Name Ataxin-10 (Brain protein E46 homolog) (Spinocerebellar ataxia type 10 lmmunogen Synthesized peptide derived from human protein . at AA range: 10-90 Specificity ATX10 Polyclonal Antibody detects endogenous levels of protein.  Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.  Source Polyclonal, Rabbit,IgG  Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000 ELISA 1:5000-20000   | protein)                         |
| Gene Name ATXN10 SCA10  Protein Name Ataxin-10 (Brain protein E46 homolog) (Spinocerebellar ataxia type 10 Immunogen Synthesized peptide derived from human protein . at AA range: 10-90 Specificity ATX10 Polyclonal Antibody detects endogenous levels of protein.  Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.  Source Polyclonal, Rabbit,IgG  Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000 ELISA 1:5000-20000  | protein)                         |
| Protein NameAtaxin-10 (Brain protein E46 homolog) (Spinocerebellar ataxia type 10ImmunogenSynthesized peptide derived from human protein . at AA range: 10-90SpecificityATX10 Polyclonal Antibody detects endogenous levels of protein.FormulationLiquid in PBS containing 50% glycerol, and 0.02% sodium azide.SourcePolyclonal, Rabbit,IgGPurificationThe antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.DilutionWB 1:500-2000 ELISA 1:5000-20000   | protein)                         |
| ImmunogenSynthesized peptide derived from human protein . at AA range: 10-90SpecificityATX10 Polyclonal Antibody detects endogenous levels of protein.FormulationLiquid in PBS containing 50% glycerol, and 0.02% sodium azide.SourcePolyclonal, Rabbit,IgGPurificationThe antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.DilutionWB 1:500-2000 ELISA 1:5000-20000  | protein)                         |
| SpecificityATX10 Polyclonal Antibody detects endogenous levels of protein.FormulationLiquid in PBS containing 50% glycerol, and 0.02% sodium azide.SourcePolyclonal, Rabbit,IgGPurificationThe antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.DilutionWB 1:500-2000 ELISA 1:5000-20000  |                                  |
| Formulation  Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.  Source  Polyclonal, Rabbit,IgG  Purification  The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution  WB 1:500-2000 ELISA 1:5000-20000  |                                  |
| Source Polyclonal, Rabbit, IgG  Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000 ELISA 1:5000-20000   |                                  |
| Purification  The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution  WB 1:500-2000 ELISA 1:5000-20000   |                                  |
| affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000 ELISA 1:5000-20000  |                                  |
|   |                                  |
| Concentration 1 mg/ml   |                                  |
|   |                                  |
| Purity ≥90%   |                                  |
| Storage Stability -20°C/1 year  |                                  |
| Synonyms  |                                  |
| Observed Band 52kD  |                                  |
| Cell Pathway Cytoplasm, perinuclear region .  |                                  |
| Tissue Specificity Expressed in the central nervous system.   |                                  |
| Function  disease:Defects in ATXN10 are the cause of spinocerebellar ataxia typ (SCA10) [MIM:603516]. Spinocerebellar ataxia is a clinically and genet heterogeneous group of cerebellar disorders. Patients show progressiv incoordination of gait and often poor coordination of hands, speech and movements, due to degeneration of the cerebellum with variable involve the brainstem and spinal cord. SCA10 is an autosomal dominant cerebe (ADCA).,similarity:Belongs to the ataxin-10 family.,   | ically<br>e<br>I eye<br>ement of |
|   | ellar ataxia                     |

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658







|                           | associated with spinocerebellar ataxia, type 10. Alternatively spliced transcript variants have been described.[provided by RefSeq, Jul 2016], |  |
|---------------------------|--|--|
| matters needing attention | Avoid repeated freezing and thawing!   |  |
| Usage suggestions         | This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.              |  |

| Products Images |
|-----------------|
|                 |
|                 |
|                 |
|                 |
|                 |