



CEAM7 Polyclonal Antibody

BYab-05435
IgG
Human;Rat;Mouse;
WB;ELISA
CEACAM7 CGM2
Carcinoembryonic antigen-related cell adhesion molecule 7 (Carcinoembryonic antigen CGM2)
Synthesized peptide derived from part region of human protein
CEAM7 Polyclonal Antibody detects endogenous levels of protein.
Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Polyclonal, Rabbit,IgG
The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
WB 1:500-2000 ELISA 1:5000-20000
1 mg/ml
≥90%
-20°C/1 year
29kD
Cell membrane ; Lipid-anchor, GPI-anchor . Apical cell membrane . Localized to the apical glycocalyx surface
Expressed in columnar epithelial cells of the colon (at protein level) (PubMed:10436421). Strongly down-regulated in colonic adenocarcinomas.
similarity:Belongs to the immunoglobulin superfamily. CEA familysimilarity:Contains 1 Ig-like C2-type (immunoglobulin-like)
domáin.,similarity:Contains 1 lg-like V-týpe (immunoglobulin-like) domain.,tissue specificity:Strongly down-regulated in colonic adenocarcinomas.,

Nanjing BYabscience technology Co.,Ltd








matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western blot analysis of lysates from A431 cells, primary antibody was diluted at 1:1000, 4° over night

Nanjing BYabscience technology Co.,Ltd