



# CTSR4 Polyclonal Antibody

|                           |  |
|---------------------------|--|
| <b>Catalog No</b>         | BYab-05440   |
| <b>Isotype</b>            | IgG  |
| <b>Reactivity</b>         | Human;Rat;Mouse;   |
| <b>Applications</b>       | WB;ELISA   |
| <b>Gene Name</b>          | CATSPER4   |
| <b>Protein Name</b>       | Cation channel sperm-associated protein 4 (CatSper4)   |
| <b>Immunogen</b>          | Synthesized peptide derived from part region of human protein  |
| <b>Specificity</b>        | CTSR4 Polyclonal Antibody detects endogenous levels of protein.  |
| <b>Formulation</b>        | Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.   |
| <b>Source</b>             | Polyclonal, Rabbit,IgG   |
| <b>Purification</b>       | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  |
| <b>Dilution</b>           | WB 1:500-2000 ELISA 1:5000-20000   |
| <b>Concentration</b>      | 1 mg/ml  |
| <b>Purity</b>             | ≥90%   |
| <b>Storage Stability</b>  | -20°C/1 year   |
| <b>Synonyms</b>           |  |
| <b>Observed Band</b>      | 51kD   |
| <b>Cell Pathway</b>       | Cell projection, cilium, flagellum membrane ; Multi-pass membrane protein . Specifically located in the principal piece of the sperm tail. .   |
| <b>Tissue Specificity</b> | Testis-specific.   |
| <b>Function</b>           | function:Voltage-gated calcium channel that plays a central role in sperm cell hyperactivation. Controls calcium entry to mediate the hyperactivated motility, a step needed for sperm motility which is essential late in the preparation of sperm for fertilization. Activated by intracellular alkalinization.,similarity:Belongs to the cation channel sperm-associated family.,subunit:Heterotetramer; possibly composed of CATSPER1, CATSPER2, CATSPER3 and CATSPER4 (Potential). Interacts with CATSPER1.,tissue specificity:Testis-specific. Specifically present in the principal piece of sperm tail., |
| <b>Background</b>         | function:Voltage-gated calcium channel that plays a central role in sperm cell hyperactivation. Controls calcium entry to mediate the hyperactivated motility, a step needed for sperm motility which is essential late in the preparation of sperm  |

Nanjing BYabscience technology Co.,Ltd



for fertilization. Activated by intracellular alkalinization.,similarity:Belongs to the cation channel sperm-associated family.,subunit:Heterotetramer; possibly composed of CATSPER1, CATSPER2, CATSPER3 and CATSPER4 (Potential). Interacts with CATSPER1.,tissue specificity:Testis-specific. Specifically present in the principal piece of sperm tail.,

**matters needing attention**

Avoid repeated freezing and thawing!

**Usage suggestions**

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

**Products Images**