



# DSG3 Polyclonal Antibody

<b>Catalog No</b>	BYab-05514
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Rat;Mouse;
<b>Applications</b>	WB;ELISA
<b>Gene Name</b>	DSG3 CDHF6
<b>Protein Name</b>	Desmoglein-3 (130 kDa pemphigus vulgaris antigen) (PVA) (Cadherin family member 6)
<b>Immunogen</b>	Synthesized peptide derived from part region of human protein
<b>Specificity</b>	DSG3 Polyclonal Antibody detects endogenous levels of protein.
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
<b>Source</b>	Polyclonal, Rabbit,IgG
<b>Purification</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Dilution</b>	WB 1:500-2000 ELISA 1:5000-20000
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	
<b>Observed Band</b>	109kD
<b>Cell Pathway</b>	Cell membrane ; Single-pass type I membrane protein . Cell junction, desmosome .
<b>Tissue Specificity</b>	Epidermis, tongue, tonsil, esophagus and carcinomas. Expressed in skin and mucosa (at protein level) (PubMed:30528827).
<b>Function</b>	disease:Pemphigus vulgaris (PV) is a potentially lethal skin disease in which epidermal blisters occur as the result of the loss of cell-cell adhesion caused by the action of autoantibodies against desmoglein 3.,domain:Calcium may be bound by the cadherin-like repeats .,function:Component of intercellular desmosome junctions. Involved in the interaction of plaque proteins and intermediate filaments mediating cell-cell adhesion.,similarity:Contains 4 cadherin domains.,tissue specificity:Epidermis, tongue, tonsil, esophagus and carcinomas.,
<b>Background</b>	This gene encodes a member of the desmoglein family and cadherin cell adhesion molecule superfamily of proteins. Desmogleins are calcium-binding transmembrane glycoprotein components of desmosomes, cell-cell junctions between epithelial, myocardial, and other cell types. The encoded preproprotein is

**Nanjing BYabscience technology Co.,Ltd**



proteolytically processed to generate the mature glycoprotein. This gene is present in a gene cluster with other desmoglein gene family members on chromosome 18. The encoded protein has been identified as the autoantigen of the autoimmune blistering disease pemphigus vulgaris. [provided by RefSeq, Jan 2016],

**matters needing attention**

Avoid repeated freezing and thawing!

**Usage suggestions**

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

**Products Images**