



## HNRH3 Polyclonal Antibody

isoforms may possess different functions in the splicing reaction.,similarity:Contains 2 RRM (RNA recognition motif) domains.,  This gene belongs to the subfamily of ubiquitously expressed heterogeneous nuclear ribonucleoproteins (hnRNPs). The hnRNPs are RNA binding proteins at they complex with heterogeneous nuclear RNA (hnRNA). These proteins are associated with pre-mRNAs in the nucleus and appear to influence pre-mRNA processing and other aspects of mRNA metabolism and transport. While all of the		
Reactivity Human;Rat;Mouse;  Applications WB;ELISA  Gene Name HNRNPH3 HNRPH3  Protein Name Heterogeneous nuclear ribonucleoprotein H3 (hnRNP H3) (Heterogeneous nuclear ribonucleoprotein 2H9) (hnRNP 2H9)  Immunogen Synthesized peptide derived from part region of human protein  Specificity HNRH3 Polyclonal Antibody detects endogenous levels of protein.  Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.  Source Polyclonal, Rabbit,IgG  Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000 ELISA 1:5000-20000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band 38kD  Cell Pathway Nucleus.  Tissue Specificity Brain,Cajal-Retzius cell,Cerebellum,Epithelium,Fetal brain  function:Involved in the splicing process and participates in early heat shock-induced splicing arrest. Due to their great structural variations the difference isoforms may possess different inctions in the splicing reaction.,similarity:Contains 2 RRM (RNA recognition motif) domains.  Background This gene belongs to the subfamily of ubiquitously expressed heterogeneous they complex with heterogeneous nuclear RNA (hnRNA). These proteins are associated with pre-mRNAs in the nucleus and appear to influence pre-mRNA processing and other aspects of mRNA metabolism and transport. Wile all of the nRNPs are RNA in fine nucleus, some seem to shuttle between the nucleus	Catalog No	BYab-05636
Applications WB;ELISA  Gene Name HNRNPH3 HNRPH3  Protein Name Heterogeneous nuclear ribonucleoprotein H3 (hnRNP H3) (Heterogeneous nuclear ribonucleoprotein 2H9) (hnRNP 2H9)  Immunogen Synthesized peptide derived from part region of human protein  Specificity HNRH3 Polyclonal Antibody detects endogenous levels of protein.  Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.  Source Polyclonal, Rabbit,IgG  Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000 ELISA 1:5000-20000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band 38kD  Cell Pathway Nucleus.  Tissue Specificity Brain, Cajal-Retzius cell, Cerebellum, Epithelium, Fetal brain function: Involved in the splicing process and participates in early heat shock-induced splicing arrest. Due to their great structural variations the different isoforms may possess different functions in the splicing reaction., similarity: Contains 2 RRM (RNA recognition motif) domains.,  This gene belongs to the subfamily of ubiquitously expressed heterogeneous nuclear ribonucleoproteins (hnRNPs). The hnRNPs are RNA binding proteins at they complex with heterogeneous nuclear RNA (hnRNA). These proteins are associated with pre-mRNAs in the nucleus and appear to influence pre-mRNA processing and transport. While all of the NRNPs are present in the nucleus, some seem to shuttle between the nucleus	Isotype	IgG
Gene Name HNRNPH3 HNRPH3  Protein Name Heterogeneous nuclear ribonucleoprotein H3 (hnRNP H3) (Heterogeneous nuclear ribonucleoprotein 2H9) (hnRNP 2H9)  Immunogen Synthesized peptide derived from part region of human protein  Specificity HNRH3 Polyclonal Antibody detects endogenous levels of protein.  Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.  Source Polyclonal, Rabbit, IgG  Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000 ELISA 1:5000-20000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band 38kD  Cell Pathway Nucleus.  Tissue Specificity Brain, Cajal-Retzius cell, Cerebellum, Epithelium, Fetal brain function: Involved in the splicing process and participates in early heat shock-induced splicing arrest. Due to their great structural variations the different isoforms may possess different functions in the splicing reaction., similarity: Contains 2 RRM (RNA recognition motif) domains.,  Background This gene belongs to the subfamily of bubiquitously expressed heterogeneous nuclear RNA (hnRNA). These proteins are associated with pre-mRNAs in the nucleus and appear to influence pre-mRNA processing and other aspects of mRNA metabolism and transport. While all tolerus.	Reactivity	Human;Rat;Mouse;
Protein Name Heterogeneous nuclear ribonucleoprotein H3 (hnRNP H3) (Heterogeneous nuclear ribonucleoprotein 2H9) (hnRNP 2H9)  Immunogen Synthesized peptide derived from part region of human protein  Specificity HNRH3 Polyclonal Antibody detects endogenous levels of protein.  Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.  Source Polyclonal, Rabbit,IgG  Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000 ELISA 1:5000-20000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band 38kD  Cell Pathway Nucleus.  Tissue Specificity Brain,Cajal-Retzius cell,Cerebellum,Epithelium,Fetal brain function:Involved in the splicing process and participates in early heat shock-induced splicing arrest. Due to their great structural variations the different isoforms may possess different functions in the splicing reaction.,similarity:Contains 2 RRM (RNA recognition motif) domains.,  This gene belongs to the subfamily of ubiquitously expressed heterogeneous nuclear ribonucleoproteins (nRNPS). The hnRNPs are RNA binding proteins are associated with pre-mRNAs in the nucleus and appear to influence pre-mRNA processing and other aspects of mRNA metabolism and transport. While all of the NRNPs are present in the nucleus, some seem to shuttle between the nucleus	Applications	WB;ELISA
Immunogen Synthesized peptide derived from part region of human protein  Specificity HNRH3 Polyclonal Antibody detects endogenous levels of protein.  Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.  Source Polyclonal, Rabbit, IgG  Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000 ELISA 1:5000-20000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band 38kD  Cell Pathway Nucleus.  Tissue Specificity Brain, Cajal-Retzius cell, Cerebellum, Epithelium, Fetal brain shock-induced splicing arrest. Due to their great structural variations the different isoforms may possess different functions in the splicing reaction, similantly. Contains 2 RRM (RNA recognition motif) domains.  This gene belongs to the subfamily of ubiquitously expressed heterogeneous nuclear RNA (hnRNA). These proteins are associated with pre-mRNAs in the nucleus and appear to influence pre-mRNA processing and other aspects of mRNA metabolism and transport. While all of the hnRNPs are present in the nucleus sone seem to shiftle between the nucleus.	Gene Name	HNRNPH3 HNRPH3
Immunogen         Synthesized peptide derived from part region of human protein           Specificity         HNRH3 Polyclonal Antibody detects endogenous levels of protein.           Formulation         Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.           Source         Polyclonal, Rabbit, IgG           Purification         The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.           Dilution         WB 1:500-2000 ELISA 1:5000-20000           Concentration         1 mg/ml           Purity         ≥90%           Storage Stability         -20°C/1 year           Synonyms           Observed Band         38kD           Cell Pathway         Nucleus .           Tissue Specificity         Brain, Cajal-Retzius cell, Cerebellum, Epithelium, Fetal brain           Function         function: Involved in the splicing process and participates in early heat shock-induced splicing arrest. Due to their great structural variations the different isoforms may possess different functions in the splicing reaction, similarity. Contains 2 RRM (RNA recognition motif) domains.,           Background         This gene belongs to the subtamily of ubiquitously expressed heterogeneous nuclear RNA (hnRNA). Thee proteins are associated with pre-mRNAs in the nucleus and appear to influence pre-mRNAs in the nucl	Protein Name	Heterogeneous nuclear ribonucleoprotein H3 (hnRNP H3) (Heterogeneous nuclear ribonucleoprotein 2H9) (hnRNP 2H9)
Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.  Source Polyclonal, Rabbit, IgG  Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000 ELISA 1:5000-20000  Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year  Synonyms  Observed Band 38kD  Cell Pathway Nucleus Tissue Specificity Brain, Cajal-Retzius cell, Cerebellum, Epithelium, Fetal brain  function: Involved in the splicing process and participates in early heat shock-induced splicing arrest. Due to their great structural variations the different isoforms may possess different functions in the splicing reaction., similarity: Contains 2 RRM (RNA recognition motif) domains.,  Background This gene belongs to the subfamily of ubiquitously expressed heterogeneous nuclear ribonucleoproteins (InRNPs). The hnRNPs are PNA binding proteins at they complex with heterogeneous nuclear RNA (hnRNA). These proteins are associated with pre-mRNAs in the nucleus and appear to influence pre-mRNA processing and other aspects of mRNA metabolism and transport. While all of the nRNPs are present in the nucleus, some seem to shuttle between the nucleus.	Immunogen	· · · · · · · · · · · · · · · · · · ·
Source       Polyclonal, Rabbit,IgG         Purification       The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.         Dilution       WB 1:500-2000 ELISA 1:5000-20000         Concentration       1 mg/ml         Purity       ≥90%         Storage Stability       -20°C/1 year         Synonyms       Observed Band       38kD         Cell Pathway       Nucleus .         Tissue Specificity       Brain, Cajal-Retzius cell, Cerebellum, Epithelium, Fetal brain         Function       function: Involved in the splicing process and participates in early heat shock-induced splicing arrest. Due to their great structural variations the different isoforms may possess different functions in the splicing reaction., similarity: Contains 2 RRM (RNA recognition motif) domains.,         Background       This gene belongs to the subfamily of ubiquitously expressed heterogeneous nuclear RNA (nRNAPs). These proteins are associated with pre-mRNAs in the nucleus and appear to influence pre-mRNA processing and other aspects of mRNA metabolism and transport. While all of the hRNPs are present in the nucleus, some seem to shuttle between the nucleus	Specificity	HNRH3 Polyclonal Antibody detects endogenous levels of protein.
Purification  The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution  WB 1:500-2000 ELISA 1:5000-20000  Concentration  1 mg/ml  Purity  ≥90%  Storage Stability  -20°C/1 year  Synonyms  Observed Band  38kD  Cell Pathway  Nucleus .  Tissue Specificity  Brain, Cajal-Retzius cell, Cerebellum, Epithelium, Fetal brain  function: Involved in the splicing process and participates in early heat shock-induced splicing arrest. Due to their great structural variations the different isoforms may possess different functions in the splicing reaction., similarity: Contains 2 RRM (RNA recognition motif) domains.,  Background  This gene belongs to the subfamily of ubiquitously expressed heterogeneous nuclear ribonucleoproteins (hnRNPs). The hnRNPs are RNA binding proteins at they complex with heterogeneous nuclear RNA (hnRNA). These proteins are associated with pre-mRNAs in the nucleus and appear to influence pre-mRNA processing and other aspects of mRNA metabolism and transport. While all of the hnRNPs are present in the nucleus, some seem to shuttle between the nucleus.	Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000 ELISA 1:5000-20000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band 38kD  Cell Pathway Nucleus.  Tissue Specificity Brain, Cajal-Retzius cell, Cerebellum, Epithelium, Fetal brain  function: Involved in the splicing process and participates in early heat shock-induced splicing arrest. Due to their great structural variations the different isoforms may possess different functions in the splicing reaction., similarity: Contains 2 RRM (RNA recognition motif) domains.,  Background This gene belongs to the subfamily of ubiquitously expressed heterogeneous and they complex with heterogeneous nuclear RNA (hnRNA). These proteins are associated with pre-mRNAs in the nucleus and appear to influence pre-mRNA processing and other aspects of mRNA metabolism and transport. While all of the nRNPs are present in the nucleus, some seem to shuttle between the nucleus.	Source	Polyclonal, Rabbit,IgG
Concentration       1 mg/ml         Purity       ≥90%         Storage Stability       -20°C/1 year         Synonyms       -20°C/1 year         Observed Band       38kD         Cell Pathway       Nucleus .         Tissue Specificity       Brain,Cajal-Retzius cell,Cerebellum,Epithelium,Fetal brain         Function       function:Involved in the splicing process and participates in early heat shock-induced splicing arrest. Due to their great structural variations the difference isoforms may possess different functions in the splicing reaction.,similarity:Contains 2 RRM (RNA recognition motif) domains.,         Background       This gene belongs to the subfamily of ubiquitously expressed heterogeneous nuclear ribonucleoproteins (hnRNPs). The hnRNPs are RNA binding proteins are associated with pre-mRNAs in the nucleus and appear to influence pre-mRNA processing and other aspects of mRNA metabolism and transport. While all of the hnRNPs are present in the nucleus, some seem to shuttle between the nucleus	Purification	· · · · · · · · · · · · · · · · · · ·
Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band 38kD  Cell Pathway Nucleus .  Tissue Specificity Brain,Cajal-Retzius cell,Cerebellum,Epithelium,Fetal brain  Function function:Involved in the splicing process and participates in early heat shock-induced splicing arrest. Due to their great structural variations the different isoforms may possess different functions in the splicing reaction.,similarity:Contains 2 RRM (RNA recognition motif) domains.,  Background This gene belongs to the subfamily of ubiquitously expressed heterogeneous nuclear ribonucleoproteins (hnRNPs). The hnRNPs are RNA binding proteins are associated with pre-mRNAs in the nucleus and appear to influence pre-mRNA processing and other aspects of mRNA metabolism and transport. While all of the hnRNPs are present in the nucleus, some seem to shuttle between the nucleus	Dilution	WB 1:500-2000 ELISA 1:5000-20000
Synonyms  Observed Band 38kD  Cell Pathway Nucleus .  Tissue Specificity Brain, Cajal-Retzius cell, Cerebellum, Epithelium, Fetal brain  Function function: Involved in the splicing process and participates in early heat shock-induced splicing arrest. Due to their great structural variations the different isoforms may possess different functions in the splicing reaction., similarity: Contains 2 RRM (RNA recognition motif) domains.,  Background This gene belongs to the subfamily of ubiquitously expressed heterogeneous nuclear ribonucleoproteins (hnRNPs). The hnRNPs are RNA binding proteins are associated with pre-mRNAs in the nucleus and appear to influence pre-mRNA processing and other aspects of mRNA metabolism and transport. While all of the hnRNPs are present in the nucleus, some seem to shuttle between the nucleus	Concentration	1 mg/ml
Synonyms  Observed Band 38kD  Cell Pathway Nucleus .  Tissue Specificity Brain, Cajal-Retzius cell, Cerebellum, Epithelium, Fetal brain  Function function: Involved in the splicing process and participates in early heat shock-induced splicing arrest. Due to their great structural variations the different isoforms may possess different functions in the splicing reaction., similarity: Contains 2 RRM (RNA recognition motif) domains.,  Background This gene belongs to the subfamily of ubiquitously expressed heterogeneous nuclear ribonucleoproteins (hnRNPs). The hnRNPs are RNA binding proteins are associated with pre-mRNAs in the nucleus and appear to influence pre-mRNA processing and other aspects of mRNA metabolism and transport. While all of the hnRNPs are present in the nucleus, some seem to shuttle between the nucleus.	Purity	≥90%
Observed Band  Cell Pathway  Nucleus .  Tissue Specificity  Brain,Cajal-Retzius cell,Cerebellum,Epithelium,Fetal brain  function:Involved in the splicing process and participates in early heat shock-induced splicing arrest. Due to their great structural variations the different isoforms may possess different functions in the splicing reaction.,similarity:Contains 2 RRM (RNA recognition motif) domains.,  Background  This gene belongs to the subfamily of ubiquitously expressed heterogeneous nuclear ribonucleoproteins (hnRNPs). The hnRNPs are RNA binding proteins are associated with pre-mRNAs in the nucleus and appear to influence pre-mRNA processing and other aspects of mRNA metabolism and transport. While all of the hnRNPs are present in the nucleus, some seem to shuttle between the nucleus	Storage Stability	-20°C/1 year
Tissue Specificity  Brain,Cajal-Retzius cell,Cerebellum,Epithelium,Fetal brain  function:Involved in the splicing process and participates in early heat shock-induced splicing arrest. Due to their great structural variations the different isoforms may possess different functions in the splicing reaction.,similarity:Contains 2 RRM (RNA recognition motif) domains.,  This gene belongs to the subfamily of ubiquitously expressed heterogeneous nuclear ribonucleoproteins (hnRNPs). The hnRNPs are RNA binding proteins are associated with pre-mRNAs in the nucleus and appear to influence pre-mRNA processing and other aspects of mRNA metabolism and transport. While all of the hnRNPs are present in the nucleus, some seem to shuttle between the nucleus	Synonyms	
Tissue Specificity  Brain, Cajal-Retzius cell, Cerebellum, Epithelium, Fetal brain  function: Involved in the splicing process and participates in early heat shock-induced splicing arrest. Due to their great structural variations the difference isoforms may possess different functions in the splicing reaction., similarity: Contains 2 RRM (RNA recognition motif) domains.,  This gene belongs to the subfamily of ubiquitously expressed heterogeneous nuclear ribonucleoproteins (hnRNPs). The hnRNPs are RNA binding proteins are associated with pre-mRNAs in the nucleus and appear to influence pre-mRNA processing and other aspects of mRNA metabolism and transport. While all of the hnRNPs are present in the nucleus, some seem to shuttle between the nucleus	Observed Band	38kD
function: Involved in the splicing process and participates in early heat shock-induced splicing arrest. Due to their great structural variations the difference isoforms may possess different functions in the splicing reaction., similarity: Contains 2 RRM (RNA recognition motif) domains.,  This gene belongs to the subfamily of ubiquitously expressed heterogeneous nuclear ribonucleoproteins (hnRNPs). The hnRNPs are RNA binding proteins are they complex with heterogeneous nuclear RNA (hnRNA). These proteins are associated with pre-mRNAs in the nucleus and appear to influence pre-mRNA processing and other aspects of mRNA metabolism and transport. While all of the hnRNPs are present in the nucleus, some seem to shuttle between the nucleus	Cell Pathway	Nucleus .
shock-induced splicing arrest. Due to their great structural variations the different isoforms may possess different functions in the splicing reaction., similarity: Contains 2 RRM (RNA recognition motif) domains.,  This gene belongs to the subfamily of ubiquitously expressed heterogeneous nuclear ribonucleoproteins (hnRNPs). The hnRNPs are RNA binding proteins at they complex with heterogeneous nuclear RNA (hnRNA). These proteins are associated with pre-mRNAs in the nucleus and appear to influence pre-mRNA processing and other aspects of mRNA metabolism and transport. While all of the hnRNPs are present in the nucleus, some seem to shuttle between the nucleus	Tissue Specificity	Brain,Cajal-Retzius cell,Cerebellum,Epithelium,Fetal brain
nuclear ribonucleoproteins (hnRNPs). The hnRNPs are RNA binding proteins and they complex with heterogeneous nuclear RNA (hnRNA). These proteins are associated with pre-mRNAs in the nucleus and appear to influence pre-mRNA processing and other aspects of mRNA metabolism and transport. While all of the hnRNPs are present in the nucleus, some seem to shuttle between the nucleus.	Function	shock-induced splicing arrest. Due to their great structural variations the different isoforms may possess different functions in the splicing
	Background	nuclear ribonucleoproteins (hnRNPs). The hnRNPs are RNA binding proteins and they complex with heterogeneous nuclear RNA (hnRNA). These proteins are associated with pre-mRNAs in the nucleus and appear to influence pre-mRNA processing and other aspects of mRNA metabolism and transport. While all of the hnRNPs are present in the nucleus, some seem to shuttle between the nucleus

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658




国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询



	properties. The protein encoded by this gene has two repeats of quasi-RRM domains that bind to RNAs. It is localized in nuclear bodies of the nucleus. This protein is involved in the splicing process and it also participates in early heat shock-induced splicing arrest by transiently leaving the hnRNP complexes. Several alternatively spliced transcript variant
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## **Products Images**



Western blot analysis of lysates from DU145 cells, primary antibody was diluted at 1:1000, 4° over night

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658