



LRC42 Polyclonal Antibody

Catalog No BYab-05690 Isotype IgG Reactivity Human;Mouse;Rat Applications WB;ELISA Gene Name LRRC42 Protein Name Leucine-rich repeat-containing protein 42 Immunogen Synthesized peptide derived from part region of human protein Specificity LRC42 Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 47kD Cell Pathway Embryo, Placenta, Skin, Function PTM:Phosphorylated upon DNA damage, probably by ATM or ATR, similarity:Belongs to the LRRC42 family, similarity:Contains 2 LRR (leucine-rich) repeats. Background Avoid repeated freezing and thawing!		
Reactivity Human;Mouse;Rat Applications WB;ELISA Gene Name LRRC42 Protein Name Leucine-rich repeat-containing protein 42 Immunogen Synthesized peptide derived from part region of human protein Specificity LRC42 Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 47kD Cell Pathway Tissue Specificity Embryo,Placenta,Skin, Function PTM:Phosphorylated upon DNA damage, probably by ATM or ATR_similarity:Belongs to the LRRC42 family_similarity:Contains 2 LRR (leucine-rich) repeats. Background Avoid repeated freezing and thawing!	Catalog No	BYab-05690
Applications WB;ELISA Gene Name LRRC42 Protein Name Leucine-rich repeat-containing protein 42 Immunogen Synthesized peptide derived from part region of human protein Specificity LRC42 Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 47kD Cell Pathway Tissue Specificity Embryo,Placenta,Skin, Function PTM:Phosphorylated upon DNA damage, probably by ATM or ATR, similarity Belongs to the LRRC42 family, similarity:Contains 2 LRR (leucine-rich) repeats. Background Avoid repeated freezing and thawing!	Isotype	IgG
Gene Name LRRC42 Protein Name Leucine-rich repeat-containing protein 42 Immunogen Synthesized peptide derived from part region of human protein Specificity LRC42 Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 47kD Cell Pathway Tissue Specificity Embryo, Placenta, Skin, Function PTM:Phosphorylated upon DNA damage, probably by ATM or ATR., similarity:Belongs to the LRRC42 family., similarity:Contains 2 LRR (leucine-rich) repeats., Background PTM:Phosphorylated upon DNA damage, probably by ATM or ATR., similarity:Belongs to the LRRC42 family., similarity:Contains 2 LRR (leucine-rich) repeats., matters needing Avoid repeated freezing and thawing!	Reactivity	Human;Mouse;Rat
Protein Name Leucine-rich repeat-containing protein 42 Immunogen Synthesized peptide derived from part region of human protein Specificity LRC42 Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 47kD Cell Pathway Embryo,Placenta,Skin, Function PTM:Phosphorylated upon DNA damage, probably by ATM or ATR, similarity:Belongs to the LRRC42 family, similarity:Contains 2 LRR (leucine-rich) repeats., Background PTM:Phosphorylated upon DNA damage, probably by ATM or ATR, similarity:Belongs to the LRRC42 family, similarity:Contains 2 LRR (leucine-rich) repeats., matters needing Avoid repeated freezing and thawing!	Applications	WB;ELISA
Immunogen Synthesized peptide derived from part region of human protein Specificity LRC42 Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 47kD Cell Pathway Tissue Specificity Embryo, Placenta, Skin, Function PTM: Phosphorylated upon DNA damage, probably by ATM or ATR., similarity: Belongs to the LRRC42 family., similarity: Contains 2 LRR (leucine-rich) repeats., Background PTM: Phosphorylated upon DNA damage, probably by ATM or ATR., similarity: Belongs to the LRRC42 family., similarity: Contains 2 LRR (leucine-rich) repeats., matters needing Avoid repeated freezing and thawing!	Gene Name	LRRC42
Specificity LRC42 Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 47kD Cell Pathway Tissue Specificity Embryo, Placenta, Skin, Function PTM: Phosphorylated upon DNA damage, probably by ATM or ATR., similarity: Belongs to the LRRC42 family., similarity: Contains 2 LRR (leucine-rich) repeats., Background PTM: Phosphorylated upon DNA damage, probably by ATM or ATR., similarity: Belongs to the LRRC42 family., similarity: Contains 2 LRR (leucine-rich) repeats., matters needing Avoid repeated freezing and thawing!	Protein Name	Leucine-rich repeat-containing protein 42
Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 47kD Cell Pathway Tissue Specificity Embryo, Placenta, Skin, Function PTM:Phosphorylated upon DNA damage, probably by ATM or ATR., similarity: Belongs to the LRRC42 family., similarity: Contains 2 LRR (leucine-rich) repeats., Background PTM:Phosphorylated upon DNA damage, probably by ATM or ATR., similarity: Belongs to the LRRC42 family., similarity: Contains 2 LRR (leucine-rich) repeats., Matters needing Avoid repeated freezing and thawing!	Immunogen	Synthesized peptide derived from part region of human protein
Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 47kD Cell Pathway Tissue Specificity Embryo, Placenta, Skin, Function PTM: Phosphorylated upon DNA damage, probably by ATM or ATR, similarity: Belongs to the LRRC42 family., similarity: Contains 2 LRR (leucine-rich) repeats., Background PTM: Phosphorylated upon DNA damage, probably by ATM or ATR, similarity: Belongs to the LRRC42 family., similarity: Contains 2 LRR (leucine-rich) repeats., Matters needing Avoid repeated freezing and thawing!	Specificity	LRC42 Polyclonal Antibody detects endogenous levels of protein.
Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 47kD Cell Pathway Tissue Specificity Embryo,Placenta,Skin, Function PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Belongs to the LRRC42 family.,similarity:Contains 2 LRR (leucine-rich) repeats., Background PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Belongs to the LRRC42 family.,similarity:Contains 2 LRR (leucine-rich) repeats., Matters needing Avoid repeated freezing and thawing!	Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 47kD Cell Pathway Tissue Specificity Embryo,Placenta,Skin, Function PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Belongs to the LRRC42 family.,similarity:Contains 2 LRR (leucine-rich) repeats., Background PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Belongs to the LRRC42 family.,similarity:Contains 2 LRR (leucine-rich) repeats., Matters needing Avoid repeated freezing and thawing!	Source	Polyclonal, Rabbit,IgG
Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 47kD Cell Pathway Tissue Specificity Embryo,Placenta,Skin, Function PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Belongs to the LRRC42 family.,similarity:Contains 2 LRR (leucine-rich) repeats., Background PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Belongs to the LRRC42 family.,similarity:Contains 2 LRR (leucine-rich) repeats., Matters needing Avoid repeated freezing and thawing!	Purification	
Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 47kD Cell Pathway Tissue Specificity Embryo,Placenta,Skin, Function PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Belongs to the LRRC42 family.,similarity:Contains 2 LRR (leucine-rich) repeats., matters needing Avoid repeated freezing and thawing!	Dilution	WB 1:500-2000 ELISA 1:5000-20000
Storage Stability -20°C/1 year Synonyms Observed Band 47kD Cell Pathway Tissue Specificity Embryo,Placenta,Skin, Function PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Belongs to the LRRC42 family.,similarity:Contains 2 LRR (leucine-rich) repeats., Background PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Belongs to the LRRC42 family.,similarity:Contains 2 LRR (leucine-rich) repeats., matters needing Avoid repeated freezing and thawing!	Concentration	1 mg/ml
Synonyms Observed Band 47kD Cell Pathway Tissue Specificity Embryo,Placenta,Skin, Function PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Belongs to the LRRC42 family.,similarity:Contains 2 LRR (leucine-rich) repeats., Background PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Belongs to the LRRC42 family.,similarity:Contains 2 LRR (leucine-rich) repeats., matters needing Avoid repeated freezing and thawing!	Purity	≥90%
Observed Band Cell Pathway Tissue Specificity Embryo,Placenta,Skin, PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Belongs to the LRRC42 family.,similarity:Contains 2 LRR (leucine-rich) repeats., Background PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Belongs to the LRRC42 family.,similarity:Contains 2 LRR (leucine-rich) repeats., matters needing Avoid repeated freezing and thawing!	Storage Stability	-20°C/1 year
Tissue Specificity Embryo,Placenta,Skin, PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Belongs to the LRRC42 family.,similarity:Contains 2 LRR (leucine-rich) repeats., Background PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Belongs to the LRRC42 family.,similarity:Contains 2 LRR (leucine-rich) repeats., matters needing Avoid repeated freezing and thawing!	Synonyms	
Tissue Specificity Embryo,Placenta,Skin, PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Belongs to the LRRC42 family.,similarity:Contains 2 LRR (leucine-rich) repeats., Background PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Belongs to the LRRC42 family.,similarity:Contains 2 LRR (leucine-rich) repeats., matters needing Avoid repeated freezing and thawing!	Observed Band	47kD
Function PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Belongs to the LRRC42 family.,similarity:Contains 2 LRR (leucine-rich) repeats., PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Belongs to the LRRC42 family.,similarity:Contains 2 LRR (leucine-rich) repeats., matters needing Avoid repeated freezing and thawing!	Cell Pathway	
ATR.,similarity:Belongs to the LRRC42 family.,similarity:Contains 2 LRR (leucine-rich) repeats., Background PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Belongs to the LRRC42 family.,similarity:Contains 2 LRR (leucine-rich) repeats., matters needing Avoid repeated freezing and thawing!	Tissue Specificity	Embryo,Placenta,Skin,
ATR.,similarity:Belongs to the LRRC42 family.,similarity:Contains 2 LRR (leucine-rich) repeats., matters needing Avoid repeated freezing and thawing!	Function	ATR.,similarity:Belongs to the LRRC42 family, similarity:Contains 2 LRR
	Background	ATR.,similarity:Belongs to the LRRC42 family.,similarity:Contains 2 LRR
	matters needing attention	Avoid repeated freezing and thawing!

Nanjing BYabscience technology Co.,Ltd




国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询



Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.



网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658