



NNTM Polyclonal Antibody

BYab-05788
IgG
Human;Mouse
WB;ELISA
NNT
NAD(P) transhydrogenase, mitochondrial (EC 1.6.1.2) (Nicotinamide nucleotide transhydrogenase) (Pyridine nucleotide transhydrogenase)
Synthesized peptide derived from human protein . at AA range: 240-320
NNTM Polyclonal Antibody detects endogenous levels of protein.
Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Polyclonal, Rabbit,IgG
The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
WB 1:500-2000 ELISA 1:5000-20000
1 mg/ml
≥90%
-20°C/1 year
119kD
Mitochondrion inner membrane ; Multi-pass membrane protein ; Matrix side .
Widely expressed with expression most readily detectable in adrenal, heart, kidney, thyroid and adipose tissues.
catalytic activity:NADPH + NAD(+) = NADP(+) + NADH.,function:The transhydrogenation between NADH and NADP is coupled to respiration and ATP
hydrolysis and functions as a proton pump across the membrane.,similarity:In the C-terminal section; belongs to the PNT beta subunit family.,similarity:In the N-terminal section; belongs to the AlaDH/PNT family.,subunit:Homodimer.,

Nanjing BYabscience technology Co.,Ltd







attention	Avoid repeated freezing and triawing:
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images