




NDUBB Polyclonal Antibody

(Complex ESSS) (CI-ESSS) (NADH-ubiquinone oxidoreductase ESSS subur (Neuronal protein 17.3) (Np17.3) (p17.3) Immunogen Synthesized peptide derived from human protein . at AA range: 70-150 Specificity NDUBB Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 16kD Cell Pathway Mitochondrion inner membrane; Single-pass membrane protein. The interactivith BCAP31 mediates mitochondria localization. Tissue Specificity Ubiquitous. Function function: Accessory subunit of the mitochondrial membrane respiratory chain NADH dehydrogenase (Complex I), that is believed to be not involved in catalyty Complex I functions in the transfer of electrons from NADH to the respiratory chain. The immediate electron acceptor for the enzyme is believed to be ubiquinone, similarity, Belongs to the complex I NDUFB11 subunit family, subunit family subunit accomplex I catalyzes the first step in the electron transport chain, the transfer of 2 electrons from NADH		
Reactivity	Catalog No	BYab-05797
Applications WB;ELISA Gene Name NDUFB11 UNQ111/PRO1064 Protein Name NADH dehydrogenase [ubiquinone] 1 beta subcomplex subunit 11, mitochondric (Complex LESSS) (CI-ESSS) (NADH-ubiquinone oxidoreductase ESSS subunit (Neuronal protein 17.3) (Np17.3) (p17.3) Immunogen Synthesized peptide derived from human protein . at AA range: 70-150 Specificity NDUBB Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 16kD Cell Pathway Mitochondrion inner membrane; Single-pass membrane protein . The interactivith BCAP31 mediates mitochondria localization. Tissue Specificity Ubiquitous. Uiquitous. Function function: Accessory subunit of the mitochondrial membrane respiratory chain. The immediate electron acceptor for the enzyme is believed to be ubiquinone, similarity:Belongs to the complex I NDUFB11 subunit family, subunit Complex I is composed of 45 different subunits, tissue specificity:Ubiquitous. NDUFB11 is a component of mitochondrial complex I. Complex I catalyzes the first step in the electron transport chain, the transfer of 2 electrons from NADH first step in the electron transport chain, the transfer of 2 electrons from NADH	Isotype	IgG
Gene Name NDUFB11 UNQ111/PRO1064 Protein Name NADH dehydrogenase [ubiquinone] 1 beta subcomplex subunit 11, mitochondric (Complex I-ESSS) (CI-ESSS) (NADH-ubiquinone oxidoreductase ESSS subunic (Neuronal protein 17.3) (Np17.3) (p17.3) Immunogen Synthesized peptide derived from human protein . at AA range: 70-150 Specificity NDUBB Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 16kD Cell Pathway Mitochondrion inner membrane ; Single-pass membrane protein . The interaction with BCAP31 mediates mitochondrial localization Tissue Specificity Ubiquitous. Function function:Accessory subunit of the mitochondrial membrane respiratory chain NADH dehydrogenase (Complex I), that is believed to be not involved in catalyz complex I unctions in the transfer of electrons from NADH to believed to be ubiquinone, similarity. Belongs to the complex I in DUFB11 subunit family	Reactivity	Human;Mouse
Protein Name NADH dehydrogenase [ubiquinone] 1 beta subcomplex subunit 11, mitochondri (Complex I-ESSS) (CI-ESSS) (NADH-lubiquinone oxidoreductase ESSS subunice (Neuronal protein 17.3) (Np17.3) (p17.3) Immunogen Synthesized peptide derived from human protein . at AA range: 70-150 Specificity NDUBB Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 16kD Cell Pathway Mitochondrion inner membrane; Single-pass membrane protein. The interaction with BCAP31 mediates mitochondria localization. Tissue Specificity Ubiquitous. function: Accessory subunit of the mitochondrial membrane respiratory chain NADH dehydrogenase (Complex I), that is believed to be not involved in catalyte Complex I functions in the transfer of electrons from NADH to the respiratory chain. The immediate electron acceptor for the enzyme shelieved to be ubiquinone, similarity: Belongs to the complex I subunit family, subunit Complex I is composed of 45 different subunits, tissue specificity: Ubiquitous. Background NDUFB11 is a component of mitochondrial complex I. Complex I catalyzes the first step in the electron transport chain, the transfer of 2 electrons from NADH	Applications	WB;ELISA
Complex ESSS (CI-ESSS) (NADH-ubiquinone oxidoreductase ESSS suburinary (Neuronal protein 17.3) (Np17.3) (p17.3)	Gene Name	NDUFB11 UNQ111/PRO1064
Specificity NDUBB Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 16kD Cell Pathway Mitochondrion inner membrane; Single-pass membrane protein. The interactivith BCAP31 mediates mitochondria localization Tissue Specificity Ubiquitous. Function function: Accessory subunit of the mitochondrial membrane respiratory chain NADH dehydrogenase (Complex I), that is believed to be not involved in cataly. Complex I functions in the transfer of electrons from NADH to the respiratory chain. The immediate electron acceptor for the enzyme is believed to be ubiquinone, similarity: Belongs to the complex I NDUFB11 subunit family, subunit: Complex I is composed of 45 different subunits., tissue specificity: Ubiquitous. Background NDUFB11 is a component of mitochondrial complex I. Complex I catalyzes the first step in the electron transport chain, the transfer of 2 electrons from NADH	Protein Name	NADH dehydrogenase [ubiquinone] 1 beta subcomplex subunit 11, mitochondrial (Complex I-ESSS) (CI-ESSS) (NADH-ubiquinone oxidoreductase ESSS subunit) (Neuronal protein 17.3) (Np17.3) (p17.3)
Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 16kD Cell Pathway Mitochondrion inner membrane; Single-pass membrane protein. The interactive with BCAP31 mediates mitochondria localization. Tissue Specificity Ubiquitous. Function function:Accessory subunit of the mitochondrial membrane respiratory chain NADH dehydrogenase (Complex I), that is believed to be not involved in catalysty Complex I functions in the transfer of electrons from NADH to the respiratory chain. The immediate electron acceptor for the enzyme is believed to be ubiquinone, similarity:Belongs to the complex I NDUFB11 subunit family., subunit:Complex I is composed of 45 different subunits, tissue specificity:Ubiquitous. Background NDUFB11 is a component of mitochondrial complex I. Complex I catalyzes the first step in the electron transport chain, the transfer of 2 electrons from NADH	lmmunogen	Synthesized peptide derived from human protein . at AA range: 70-150
Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 16kD Cell Pathway Mitochondrion inner membrane; Single-pass membrane protein. The interaction with BCAP31 mediates mitochondrial localization. Tissue Specificity Ubiquitous. Function function:Accessory subunit of the mitochondrial membrane respiratory chain NADH dehydrogenase (Complex I), that is believed to be not involved in catalys Complex I functions in the transfer of electrons from NADH to the respiratory chain. The immediate electron acceptor for the enzyme is believed to be ubiquinone, similarity; Belongs to the complex I NDUFB11 subunit family, subunit:Complex I is composed of 45 different subunits., tissue specificity:Ubiquitous, Background NDUFB11 is a component of mitochondrial complex I. Complex I catalyzes the first step in the electron transport chain, the transfer of 2 electrons from NADH	Specificity	NDUBB Polyclonal Antibody detects endogenous levels of protein.
Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 16kD Cell Pathway Mitochondrion inner membrane; Single-pass membrane protein. The interactivith BCAP31 mediates mitochondria localization Tissue Specificity Ubiquitous. Function function:Accessory subunit of the mitochondrial membrane respiratory chain NADH dehydrogenase (Complex I), that is believed to be not involved in catalystomplex I functions in the transfer of electrons from NADH to the respiratory chain. The immediate electron acceptor for the enzyme is believed to be ubiquinone, similarity. Belongs to the complex I NDUFB11 subunit family, subunit:Complex I is composed of 45 different subunits., tissue specificity: Ubiquitous. Background NDUFB11 is a component of mitochondrial complex I. Complex I catalyzes the first step in the electron transport chain, the transfer of 2 electrons from NADH	Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 16kD Cell Pathway Mitochondrion inner membrane; Single-pass membrane protein. The interactivith BCAP31 mediates mitochondria localization. Tissue Specificity Ubiquitous. Function function:Accessory subunit of the mitochondrial membrane respiratory chain NADH dehydrogenase (Complex I), that is believed to be not involved in catalyst Complex I functions in the transfer of electrons from NADH to the respiratory chain. The immediate electron acceptor for the enzyme is believed to be ubiquinone.,similarity:Belongs to the complex I NDUFB11 subunit familysubunit.Complex I is composed of 45 different subunits.,tissue specificity:Ubiquitous. Background NDUFB11 is a component of mitochondrial complex I. Complex I catalyzes the first step in the electron transport chain, the transfer of 2 electrons from NADH	Source	Polyclonal, Rabbit,IgG
Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Mitochondrion inner membrane; Single-pass membrane protein. The interactivith BCAP31 mediates mitochondria localization. Tissue Specificity Ubiquitous. Function function:Accessory subunit of the mitochondrial membrane respiratory chain NADH dehydrogenase (Complex I), that is believed to be not involved in catalystomain. The immediate electron acceptor for the enzyme is believed to be ubiquinone., similarity:Belongs to the complex I NDUFB11 subunit family., subunit:Complex I is composed of 45 different subunits., tissue specificity:Ubiquitous., Background NDUFB11 is a component of mitochondrial complex I. Complex I catalyzes the first step in the electron transport chain, the transfer of 2 electrons from NADH	Purification	·
Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 16kD Cell Pathway Mitochondrion inner membrane; Single-pass membrane protein. The interactive with BCAP31 mediates mitochondria localization. Tissue Specificity Ubiquitous. Function function:Accessory subunit of the mitochondrial membrane respiratory chain NADH dehydrogenase (Complex I), that is believed to be not involved in catalyst Complex I functions in the transfer of electrons from NADH to the respiratory chain. The immediate electron acceptor for the enzyme is believed to be ubiquinone.,similarity:Belongs to the complex I NDUFB11 subunit family.,subunit:Complex I is composed of 45 different subunits.,tissue specificity:Ubiquitous., Background NDUFB11 is a component of mitochondrial complex I. Complex I catalyzes the first step in the electron transport chain, the transfer of 2 electrons from NADH	Dilution	WB 1:500-2000 ELISA 1:5000-20000
Storage Stability -20°C/1 year Synonyms Observed Band 16kD Cell Pathway Mitochondrion inner membrane; Single-pass membrane protein. The interactive with BCAP31 mediates mitochondria localization. Tissue Specificity Ubiquitous. Function function:Accessory subunit of the mitochondrial membrane respiratory chain NADH dehydrogenase (Complex I), that is believed to be not involved in catalyth Complex I functions in the transfer of electrons from NADH to the respiratory chain. The immediate electron acceptor for the enzyme is believed to be specificity:Ubiquitous. Background NDUFB11 is a component of mitochondrial complex I. Complex I catalyzes the first step in the electron transport chain, the transfer of 2 electrons from NADH	Concentration	1 mg/ml
Synonyms Observed Band 16kD Cell Pathway Mitochondrion inner membrane; Single-pass membrane protein. The interaction with BCAP31 mediates mitochondria localization. Tissue Specificity Ubiquitous. Function function:Accessory subunit of the mitochondrial membrane respiratory chain NADH dehydrogenase (Complex I), that is believed to be not involved in catalyth Complex I functions in the transfer of electrons from NADH to the respiratory chain. The immediate electron acceptor for the enzyme is believed to be ubiquinone., similarity:Belongs to the complex I NDUFB11 subunit family., subunit:Complex I is composed of 45 different subunits., tissue specificity:Ubiquitous., NDUFB11 is a component of mitochondrial complex I. Complex I catalyzes the first step in the electron transport chain, the transfer of 2 electrons from NADH	Purity	≥90%
Observed Band Cell Pathway Mitochondrion inner membrane; Single-pass membrane protein. The interactive with BCAP31 mediates mitochondria localization. Tissue Specificity Ubiquitous. Function function:Accessory subunit of the mitochondrial membrane respiratory chain NADH dehydrogenase (Complex I), that is believed to be not involved in catalyst Complex I functions in the transfer of electrons from NADH to the respiratory chain. The immediate electron acceptor for the enzyme is believed to be ubiquinone.,similarity:Belongs to the complex I NDUFB11 subunit family.,subunit:Complex I is composed of 45 different subunits.,tissue specificity:Ubiquitous., NDUFB11 is a component of mitochondrial complex I. Complex I catalyzes the first step in the electron transport chain, the transfer of 2 electrons from NADH	Storage Stability	-20°C/1 year
Cell Pathway Mitochondrion inner membrane; Single-pass membrane protein. The interaction with BCAP31 mediates mitochondria localization. Tissue Specificity Ubiquitous. Function function:Accessory subunit of the mitochondrial membrane respiratory chain NADH dehydrogenase (Complex I), that is believed to be not involved in catalyst Complex I functions in the transfer of electrons from NADH to the respiratory chain. The immediate electron acceptor for the enzyme is believed to be ubiquinone.,similarity:Belongs to the complex I NDUFB11 subunit family.,subunit:Complex I is composed of 45 different subunits.,tissue specificity:Ubiquitous., NDUFB11 is a component of mitochondrial complex I. Complex I catalyzes the first step in the electron transport chain, the transfer of 2 electrons from NADH	Synonyms	
with BCAP31 mediates mitochondria localization. Tissue Specificity Ubiquitous. Function function:Accessory subunit of the mitochondrial membrane respiratory chain NADH dehydrogenase (Complex I), that is believed to be not involved in catalyst Complex I functions in the transfer of electrons from NADH to the respiratory chain. The immediate electron acceptor for the enzyme is believed to be ubiquinone.,similarity:Belongs to the complex I NDUFB11 subunit family.,subunit:Complex I is composed of 45 different subunits.,tissue specificity:Ubiquitous., NDUFB11 is a component of mitochondrial complex I. Complex I catalyzes the first step in the electron transport chain, the transfer of 2 electrons from NADH	Observed Band	16kD
Function function:Accessory subunit of the mitochondrial membrane respiratory chain NADH dehydrogenase (Complex I), that is believed to be not involved in catalyst Complex I functions in the transfer of electrons from NADH to the respiratory chain. The immediate electron acceptor for the enzyme is believed to be ubiquinone.,similarity:Belongs to the complex I NDUFB11 subunit family.,subunit:Complex I is composed of 45 different subunits.,tissue specificity:Ubiquitous., NDUFB11 is a component of mitochondrial complex I. Complex I catalyzes the first step in the electron transport chain, the transfer of 2 electrons from NADH	Cell Pathway	Mitochondrion inner membrane ; Single-pass membrane protein . The interaction with BCAP31 mediates mitochondria localization
NADH dehydrogenase (Complex I), that is believed to be not involved in catalyst Complex I functions in the transfer of electrons from NADH to the respiratory chain. The immediate electron acceptor for the enzyme is believed to be ubiquinone., similarity:Belongs to the complex I NDUFB11 subunit family., subunit:Complex I is composed of 45 different subunits., tissue specificity:Ubiquitous., NDUFB11 is a component of mitochondrial complex I. Complex I catalyzes the first step in the electron transport chain, the transfer of 2 electrons from NADH	Tissue Specificity	Ubiquitous.
first step in the electron transport chain, the transfer of 2 electrons from NADH	Function	NADH dehydrogenase (Complex I), that is believed to be not involved in catalysis. Complex I functions in the transfer of electrons from NADH to the respiratory chain. The immediate electron acceptor for the enzyme is believed to be ubiquinone., similarity:Belongs to the complex I NDUFB11 subunit family., subunit:Complex I is composed of 45 different subunits., tissue
ubiquinone, coupled to the translocation of 4 protons across the membrane	Background	NDUFB11 is a component of mitochondrial complex I. Complex I catalyzes the first step in the electron transport chain, the transfer of 2 electrons from NADH to ubiquinone, coupled to the translocation of 4 protons across the membrane

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658







(Carroll et al., 2002 [PubMed 12381726]).[supplied by OMIM, Feb 2009],

matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

