




NETO2 Polyclonal Antibody

Catalog No BYab-05853 Isotype IgG Reactivity Human;Rat;Mouse Applications WB;ELISA Gene Name NETO2 BTC12 UNQ1926/PRO4401 Protein Name Neuropilin and tolloid-like protein 2 (Brain-specific transmembrane protein containing 2 CUB and 1 LDL-receptor class A domains protein 2) Immunogen Synthesized peptide derived from human protein . at AA range: 140-220 Specificity NETO2 Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 57kD Cell Pathway Membrane; Single-pass type I membrane protein Insulative Contains 2 CUB domains followed by a low-density lipoprotein containing two circuitry, similarity:Contains 1 LDL-receptor class A domain, similarity:Contains 2 CUB domains, in the brain by regulating kainate receptor function. Expression of this gene ma		
Reactivity Human;Rat;Mouse Applications WB;ELISA Gene Name NETO2 BTCL2 UNQ1926/PRO4401 Protein Name Neuropilin and tolloid-like protein 2 (Brain-specific transmembrane protein containing 2 CUB and 1 LDL-receptor class A domains protein 2) Immunogen Synthesized peptide derived from human protein . at AA range: 140-220 Specificity NETO2 Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 57kD Cell Pathway Membrane ; Single-pass type I membrane protein . Tissue Specificity Brain,Ovary,Skin, Function function:May play a role in the development and/or maintenance of neuronal circuitry, similarity:Contains 1 LDL-receptor class A domain.,similarity:Contains 2 CUB domains. This gene encodes a predicted transmembrane protein containing two extracellular CUB domains followed by a low-density lipoprotein class A (LDLa) domains. A similar gene in rats encodes a protein that modular leases A (LDLa) domains. A similar gene in rats encodes a protein that modular leases A (LDLa) domain. A similar gene in rats encodes a protein that modular leases A (LDLa) domain. Supen in the protein protein from the development of this gene is located on the long arm of chromosome 8. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene.	Catalog No	BYab-05853
Applications WB;ELISA Gene Name NETO2 BTCL2 UNQ1926/PRO4401 Protein Name Neuropilin and tolloid-like protein 2 (Brain-specific transmembrane protein containing 2 CUB and 1 LDL-receptor class A domains protein 2) Immunogen Synthesized peptide derived from human protein at AA range: 140-220 Specificity NETO2 Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 57kD Cell Pathway Membrane; Single-pass type I membrane protein . Tissue Specificity Brain, Ovary, Skin, Function function:May play a role in the development and/or maintenance of neuronal circuitry, similarity:Contains 1 LDL-receptor class A domain., similarity:Contains 2 CUB domains. Background This gene encodes a predicted transmembrane protein containing two extracellular CUB domains followed by a low-density lipoprotein class A (LDLa) domain. A similar gene in rats encodes a protein that modulates glutamate signaling in the brain by regulating kainate receptor function. Expression of this gene in stocated on the long arm of chromosome 8. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene in stocated on the long arm of chromosome 8. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene.	Isotype	IgG
Gene Name NETO2 BTCL2 UNQ1926/PRO4401 Protein Name Neuropilin and tolloid-like protein 2 (Brain-specific transmembrane protein containing 2 CUB and 1 LDL-receptor class A domains protein 2) Immunogen Synthesized peptide derived from human protein . at AA range: 140-220 Specificity NETO2 Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 57kD Cell Pathway Membrane; Single-pass type I membrane protein Tissue Specificity Function function:May play a role in the development and/or maintenance of neuronal circuitry, similarity:Contains 1 LDL-receptor class A domain., similarity:Contains 2 CUB domains. Background This gene encodes a predicted transmembrane protein containing two extracellular CUB domains followed by a low-density lipoprotein class A (LDLa) domain. A similar gene in rats encodes a protein that modulates glutamate signaling in the bra	Reactivity	Human;Rat;Mouse
Protein Name Neuropilin and tolloid-like protein 2 (Brain-specific transmembrane protein containing 2 CUB and 1 LDL-receptor class A domains protein 2) Immunogen Synthesized peptide derived from human protein . at AA range: 140-220 Specificity NETO2 Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 57kD Cell Pathway Membrane ; Single-pass type I membrane protein. Function function:May play a role in the development and/or maintenance of neuronal circuitry, similarity:Contains 1 LDL-receptor class A domain, similarity:Contains 2 CUB domains. Background This gene encodes a predicted transmembrane protein containing two extracellular CUB domains followed by a low-density lipoprotein class A (LDLa) domain. A similar gene in rats encodes a protein that modulates glutamate signaling in the brain by regulating kainate receptor function. Expression of this gene is located on the long arm of chromosome 8. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene.	Applications	WB;ELISA
Immunogen Synthesized peptide derived from human protein . at AA range: 140-220 Specificity NETO2 Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 57kD Cell Pathway Membrane; Single-pass type I membrane protein Tissue Specificity Function function:May play a role in the development and/or maintenance of neuronal circuitry, similarity:Contains 1 LDL-receptor class A domain., similarity:Contains 2 CUB domains. Background This gene encodes a predicted transmembrane protein containing two extracellular CUB domains followed by a low-density lipoprotein class A (LDLa) domain. A similar gene in rats encodes a protein that modulates glutamate signaling in the brain by regulating kainate receptor function. Expression of this gene may be a blomarker for proliferating infantile hemangiomas. A pseudogene of this gene els located on the long arm of chromosome 8. Alternatively for this gene.	Gene Name	NETO2 BTCL2 UNQ1926/PRO4401
Specificity NETO2 Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Membrane; Single-pass type I membrane protein Tissue Specificity Brain,Ovary,Skin, Function function:May play a role in the development and/or maintenance of neuronal circuitry.;similarity:Contains 1 LDL-receptor class A domain.,similarity:Contains 2 CUB domains. Background This gene encodes a predicted transmembrane protein containing two extracellular CUB domains followed by a low-density lipoprotein class A (LDLa) domain. A similar gene in rats encodes a protein that modulates glutamate signaling in the brain by regulating kainate receptor function. Expression of this gene may be a biomarker for proliferating infantile hemangiomas. A pseudogene of this gene is located on the long arm of chromosome 8. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene.	Protein Name	Neuropilin and tolloid-like protein 2 (Brain-specific transmembrane protein containing 2 CUB and 1 LDL-receptor class A domains protein 2)
Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 57kD Cell Pathway Membrane; Single-pass type I membrane protein. Tissue Specificity Brain,Ovary,Skin, Function function:May play a role in the development and/or maintenance of neuronal circuitry, similarity:Contains 1 LDL-receptor class A domain.,similarity:Contains 2 CUB domains. Background This gene encodes a predicted transmembrane protein containing two extracellular CUB domains followed by a low-density lipoprotein class A (LDLa) domain. A similar gene in rats encodes a protein that modulates glutamate signaling in the brain by regulating kainate receptor function. Expression of this gene may be a biomarker for proliferating infantile hemangiomas. A pseudogene of this gene is located on the long arm of chromosome 8. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene.	Immunogen	Synthesized peptide derived from human protein . at AA range: 140-220
Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 57kD Cell Pathway Membrane ; Single-pass type I membrane protein . Tissue Specificity Brain,Ovary,Skin, Function function:May play a role in the development and/or maintenance of neuronal circuitry, similarity:Contains 1 LDL-receptor class A domain, similarity:Contains 2 CUB domains. Background This gene encodes a predicted transmembrane protein containing two extracellular CUB domains followed by a low-density lipoprotein class A (LDLa) domain. A similar gene in rats encodes a protein that modulates glutamate signaling in the brain by regulating kainate receptor function. Expression of this gene may be a biomarker for proliferating infantile hemangiomas. A pseudogene of this gene is located on the long arm of chromosome 8. Alternatively spliced transcript variants encoding multiple jesforms have been observed for this gene.	Specificity	NETO2 Polyclonal Antibody detects endogenous levels of protein.
Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 57kD Cell Pathway Membrane; Single-pass type I membrane protein. Tissue Specificity Brain,Ovary,Skin, Function function:May play a role in the development and/or maintenance of neuronal circuitry,similarity:Contains 1 LDL-receptor class A domain.,similarity:Contains 2 CUB domains. Background This gene encodes a predicted transmembrane protein containing two extracellular CUB domains followed by a low-density lipoprotein class A (LDLa) domain. A similar gene in rats encodes a protein that modulates glutamate signaling in the brain by regulating kainate receptor function. Expression of this gene may be a biomarker for proliferating infantile hemangiomas. A pseudogene of this gene is located on the long arm of chromosome 8. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene.	Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 57kD Cell Pathway Membrane ; Single-pass type I membrane protein . Tissue Specificity Brain,Ovary,Skin, Function function:May play a role in the development and/or maintenance of neuronal circuitry, similarity:Contains 1 LDL-receptor class A domain.,similarity:Contains 2 CUB domains. A similar gene in rats encodes a protein that modulates glutamate signaling in the brain by regulating kainate receptor function. Expression of this gene may be a biomarker for proliferating infantile hemangiomas. A pseudogene of this gene is located on the long arm of chromosome 8. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene.	Source	Polyclonal, Rabbit,IgG
Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms -20°C/1 year Observed Band 57kD Cell Pathway Membrane ; Single-pass type I membrane protein . Tissue Specificity Brain,Ovary,Skin, Function function:May play a role in the development and/or maintenance of neuronal circuitry.,similarity:Contains 1 LDL-receptor class A domain.,similarity:Contains 2 CUB domains., Background This gene encodes a predicted transmembrane protein containing two extracellular CUB domains followed by a low-density lipoprotein class A (LDLa) domain. A similar gene in rats encodes a protein that modulates glutamate signaling in the brain by regulating kainate receptor function. Expression of this gene may be a biomarker for proliferating infantile hemangiomas. A pseudogene of this gene is located on the long arm of chromosome 8. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene.	Purification	
Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 57kD Cell Pathway Membrane; Single-pass type I membrane protein. Tissue Specificity Brain,Ovary,Skin, Function function:May play a role in the development and/or maintenance of neuronal circuitry.,similarity:Contains 1 LDL-receptor class A domain.,similarity:Contains 2 CUB domains., Background This gene encodes a predicted transmembrane protein containing two extracellular CUB domains followed by a low-density lipoprotein class A (LDLa) domain. A similar gene in rats encodes a protein that modulates glutamate signaling in the brain by regulating kainate receptor function. Expression of this gene may be a biomarker for proliferating infantile hemangiomas. A pseudogene of this gene is located on the long arm of chromosome 8. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene.	Dilution	WB 1:500-2000 ELISA 1:5000-20000
Synonyms Observed Band 57kD Cell Pathway Membrane; Single-pass type I membrane protein. Tissue Specificity Brain,Ovary,Skin, Function function:May play a role in the development and/or maintenance of neuronal circuitry.,similarity:Contains 1 LDL-receptor class A domain.,similarity:Contains 2 CUB domains., Background This gene encodes a predicted transmembrane protein containing two extracellular CUB domains followed by a low-density lipoprotein class A (LDLa) domain. A similar gene in rats encodes a protein that modulates glutamate signaling in the brain by regulating kainate receptor function. Expression of this gene may be a biomarker for proliferating infantile hemangiomas. A pseudogene of this gene is located on the long arm of chromosome 8. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene.	Concentration	1 mg/ml
Synonyms Observed Band 57kD Cell Pathway Membrane; Single-pass type I membrane protein. Tissue Specificity Brain,Ovary,Skin, Function function:May play a role in the development and/or maintenance of neuronal circuitry.,similarity:Contains 1 LDL-receptor class A domain.,similarity:Contains 2 CUB domains., Background This gene encodes a predicted transmembrane protein containing two extracellular CUB domains followed by a low-density lipoprotein class A (LDLa) domain. A similar gene in rats encodes a protein that modulates glutamate signaling in the brain by regulating kainate receptor function. Expression of this gene may be a biomarker for proliferating infantile hemangiomas. A pseudogene of this gene is located on the long arm of chromosome 8. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene.	Purity	≥90%
Observed Band 57kD Cell Pathway Membrane; Single-pass type I membrane protein. Tissue Specificity Brain,Ovary,Skin, Function function:May play a role in the development and/or maintenance of neuronal circuitry.,similarity:Contains 1 LDL-receptor class A domain.,similarity:Contains 2 CUB domains., Background This gene encodes a predicted transmembrane protein containing two extracellular CUB domains followed by a low-density lipoprotein class A (LDLa) domain. A similar gene in rats encodes a protein that modulates glutamate signaling in the brain by regulating kainate receptor function. Expression of this gene may be a biomarker for proliferating infantile hemangiomas. A pseudogene of this gene is located on the long arm of chromosome 8. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene.	Storage Stability	-20°C/1 year
Cell Pathway Membrane; Single-pass type I membrane protein. Tissue Specificity Brain,Ovary,Skin, function:May play a role in the development and/or maintenance of neuronal circuitry.,similarity:Contains 1 LDL-receptor class A domain.,similarity:Contains 2 CUB domains., Background This gene encodes a predicted transmembrane protein containing two extracellular CUB domains followed by a low-density lipoprotein class A (LDLa) domain. A similar gene in rats encodes a protein that modulates glutamate signaling in the brain by regulating kainate receptor function. Expression of this gene may be a biomarker for proliferating infantile hemangiomas. A pseudogene of this gene is located on the long arm of chromosome 8. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene.	Synonyms	
Tissue Specificity Brain,Ovary,Skin, function:May play a role in the development and/or maintenance of neuronal circuitry.,similarity:Contains 1 LDL-receptor class A domain.,similarity:Contains 2 CUB domains., Background This gene encodes a predicted transmembrane protein containing two extracellular CUB domains followed by a low-density lipoprotein class A (LDLa) domain. A similar gene in rats encodes a protein that modulates glutamate signaling in the brain by regulating kainate receptor function. Expression of this gene may be a biomarker for proliferating infantile hemangiomas. A pseudogene of this gene is located on the long arm of chromosome 8. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene.	Observed Band	57kD
Function function:May play a role in the development and/or maintenance of neuronal circuitry.,similarity:Contains 1 LDL-receptor class A domain.,similarity:Contains 2 CUB domains., This gene encodes a predicted transmembrane protein containing two extracellular CUB domains followed by a low-density lipoprotein class A (LDLa) domain. A similar gene in rats encodes a protein that modulates glutamate signaling in the brain by regulating kainate receptor function. Expression of this gene may be a biomarker for proliferating infantile hemangiomas. A pseudogene of this gene is located on the long arm of chromosome 8. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene.	Cell Pathway	Membrane ; Single-pass type I membrane protein .
circuitry.,similarity:Contains 1 LDL-receptor class A domain.,similarity:Contains 2 CUB domains., Background This gene encodes a predicted transmembrane protein containing two extracellular CUB domains followed by a low-density lipoprotein class A (LDLa) domain. A similar gene in rats encodes a protein that modulates glutamate signaling in the brain by regulating kainate receptor function. Expression of this gene may be a biomarker for proliferating infantile hemangiomas. A pseudogene of this gene is located on the long arm of chromosome 8. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene.	Tissue Specificity	Brain,Ovary,Skin,
extracellular CUB domains followed by a low-density lipoprotein class A (LDLa) domain. A similar gene in rats encodes a protein that modulates glutamate signaling in the brain by regulating kainate receptor function. Expression of this gene may be a biomarker for proliferating infantile hemangiomas. A pseudogene of this gene is located on the long arm of chromosome 8. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene.	Function	circuitry.,similarity:Contains 1 LDL-receptor class A domain.,similarity:Contains 2
	Background	extracellular CUB domains followed by a low-density lipoprotein class A (LDLa)

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658







attention	Avoid repeated freezing and triawing:
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images