



NOG2 Polyclonal Antibody

| | |
|---------------------------|--|
| Catalog No | BYab-05879 |
| Isotype | IgG |
| Reactivity | Human;Mouse |
| Applications | WB;ELISA |
| Gene Name | GNL2 NGP1 |
| Protein Name | Nucleolar GTP-binding protein 2 (Autoantigen NGP-1) |
| Immunogen | Synthesized peptide derived from human protein . at AA range: 670-750 |
| Specificity | NOG2 Polyclonal Antibody detects endogenous levels of protein. |
| Formulation | Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. |
| Source | Polyclonal, Rabbit,IgG |
| Purification | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. |
| Dilution | WB 1:500-2000 ELISA 1:5000-20000 |
| Concentration | 1 mg/ml |
| Purity | ≥90% |
| Storage Stability | -20°C/1 year |
| Synonyms | |
| Observed Band | 80kD |
| Cell Pathway | Nucleus, nucleolus . |
| Tissue Specificity | Widely expressed, with the highest expression level in testis. |
| Function | function:GTPase that associates with pre-60S ribosomal subunits in the nucleolus and is required for their nuclear export and maturation.,similarity:Belongs to the MMR1/HSR1 GTP-binding protein family. NOG2 subfamily.,tissue specificity:Ubiquitously expressed at relatively low levels in all human tissues tested, with the highest level of expression in the testes., |
| Background | function:GTPase that associates with pre-60S ribosomal subunits in the nucleolus and is required for their nuclear export and maturation.,similarity:Belongs to the MMR1/HSR1 GTP-binding protein family. NOG2 subfamily.,tissue specificity:Ubiquitously expressed at relatively low levels in all human tissues tested, with the highest level of expression in the testes., |

Nanjing BYabscience technology Co.,Ltd



matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images