



PRRX2 Polyclonal Antibody

Catalog No	BYab-05896
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB;ELISA
Gene Name	PRRX2 PMX2 PRX2
Protein Name	Paired mesoderm homeobox protein 2 (Paired-related homeobox protein 2) (PRX-2)
Immunogen	Synthesized peptide derived from human protein . at AA range: 20-100
Specificity	PRRX2 Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	27kD
Cell Pathway	Nucleus .
Tissue Specificity	In fetal skin, highest expression found in cells of mesodermal origin within the dermal papilla of the developing hair shaft. Not detected in epidermis or dermis. In adult skin, weakly expressed within the basal layers of the epidermis. Not expressed in dermis.
Function	developmental stage:Higher expression in fetus than in adult.,function:May play a role in the scarless healing of cutaneous wounds during the first two trimesters of development.,similarity:Belongs to the paired homeobox family.,similarity:Contains 1 homeobox DNA-binding domain.,similarity:Contains 1 OAR domain.,tissue specificity:In fetal skin, highest expression found in cells of mesodermal origin within the dermal papilla of the developing hair shaft. Not detected in epidermis or dermis. In adult skin, weakly expressed within the basal layers of the epidermis. Not expressed in dermis.,

Nanjing BYabscience technology Co.,Ltd



Background

The DNA-associated protein encoded by this gene is a member of the paired family of homeobox proteins. Expression is localized to proliferating fetal fibroblasts and the developing dermal layer, with downregulated expression in adult skin. Increases in expression of this gene during fetal but not adult wound healing suggest a possible role in mechanisms that control mammalian dermal regeneration and prevent formation of scar response to wounding. The expression patterns provide evidence consistent with a role in fetal skin development and a possible role in cellular proliferation. [provided by RefSeq, Jul 2008],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western blot analysis of lysates from K562 cells, primary antibody was diluted at 1:1000, 4° over night