



TYSD1 Polyclonal Antibody

Catalog No	BYab-05904	
Isotype	IgG	
Reactivity	Human;Mouse	
Applications	WB;ELISA	
Gene Name	TYSND1	
Protein Name	Peroxisomal leader peptide-processing protease (EC 3.4.21) (Trypsin domain-containing protein 1) [Cleaved into: Peroxisomal leader peptide-processing protease, 15 kDa form; Peroxisomal leader peptid	
Immunogen	Synthesized peptide derived from human protein . at AA range: 120-200	
Specificity	TYSD1 Polyclonal Antibody detects endogenous levels of protein.	
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.	
Source	Polyclonal, Rabbit,IgG	
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.	
Dilution	WB 1:500-2000 ELISA 1:5000-20000	
Concentration	1 mg/ml	
Purity	≥90%	
Storage Stability	-20°C/1 year	
Synonyms		
Observed Band	62kD	
Cell Pathway	Peroxisome .	
Tissue Specificity	Brain,Lung,	
Function	function:Peroxisomal protease that mediates both the removal of the leader peptide from proteins containing a PTS2 target sequence and the specific processing of PTS1 proteins. May participate to a regulatory mechanism that control the peroxisomal beta-oxidation of fatty acids.,PTM:Processed into the 49 kDa and the 10 kDa forms upon import into the peroxisomes. The 49 kDa form probably corresponds to the mature enzyme, while the 10 kDa form may represent an inhibitory N-terminal fragment which prevents protease activity until it reaches the peroxisomes.,similarity:Belongs to the peptidase S1B family.,	
Background	This gene encodes a protease that removes the N-terminal peroxisomal targeting signal (PTS2) from proteins produced in the cytosol, thereby facilitating	

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658



国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询



	their import into the peroxisome. The encoded protein is also capable of removing the C-terminal peroxisomal targeting signal (PTS1) from proteins in the peroxisomal matrix. The full-length protein undergoes self-cleavage to produce shorter, potentially inactive, peptides. Alternative splicing results in multiple transcript variants for this gene. [provided by RefSeq, Jan 2013],
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images