



# PACS1 Polyclonal Antibody

<b>Catalog No</b>	BYab-05925
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Rat;Mouse
<b>Applications</b>	WB;ELISA
<b>Gene Name</b>	PACS1 KIAA1175
<b>Protein Name</b>	Phosphofurin acidic cluster sorting protein 1 (PACS-1)
<b>Immunogen</b>	Synthesized peptide derived from human protein . at AA range: 400-480
<b>Specificity</b>	PACS1 Polyclonal Antibody detects endogenous levels of protein.
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
<b>Source</b>	Polyclonal, Rabbit,IgG
<b>Purification</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Dilution</b>	WB 1:500-2000 ELISA 1:5000-20000
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	
<b>Observed Band</b>	105kD
<b>Cell Pathway</b>	Golgi apparatus, trans-Golgi network . Localizes in the perinuclear region, probably the TGN. .
<b>Tissue Specificity</b>	Brain,Embryo,Epithelium,Fetal brain,Kidney,Testis,Uterus,
<b>Function</b>	function:Coat protein that is involved in the localization of trans-Golgi network (TGN) membrane proteins that contain acidic cluster sorting motifs. Controls the endosome-to-Golgi trafficking of furin and mannose-6-phosphate receptor by connecting the acidic-cluster-containing cytoplasmic domain of these molecules with the adapter-protein complex-1 (AP-1) of endosomal clathrin-coated membrane pits. Involved in HIV-1 nef-mediated removal of MHC-I from the cell surface to the TGN.,PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Belongs to the PACS family.,subcellular location:Localizes in the perinuclear region, probably the TGN.,subunit:Interacts with HIV-1 Nef, AP-1 and AP-3 but not with AP-2. Forms a ternary complex with furin and AP-1. Combines with Nef to usurp the ARF6 endocytic pathway by a PI3K-dependent process to remove cell surface MHC-I molecules.,

**Nanjing BYabscience technology Co.,Ltd**



**Background**

This gene encodes a protein with a putative role in the localization of trans-Golgi network (TGN) membrane proteins. Mouse and rat homologs have been identified and studies of the homologous rat protein indicate a role in directing TGN localization of furin by binding to the protease's phosphorylated cytosolic domain. In addition, the human protein plays a role in HIV-1 Nef-mediated downregulation of cell surface MHC-I molecules to the TGN, thereby enabling HIV-1 to escape immune surveillance. [provided by RefSeq, Jul 2008],

**matters needing attention**

Avoid repeated freezing and thawing!

**Usage suggestions**

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

**Products Images**