



# PLXB1 Polyclonal Antibody

<b>Catalog No</b>	BYab-05941
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Mouse
<b>Applications</b>	WB;ELISA
<b>Gene Name</b>	PLXNB1 KIAA0407 PLXN5 SEP
<b>Protein Name</b>	Plexin-B1 (Semaphorin receptor SEP)
<b>Immunogen</b>	Synthesized peptide derived from human protein . at AA range: 90-170
<b>Specificity</b>	PLXB1 Polyclonal Antibody detects endogenous levels of protein.
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
<b>Source</b>	Polyclonal, Rabbit,IgG
<b>Purification</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Dilution</b>	WB 1:500-2000 ELISA 1:5000-20000
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	
<b>Observed Band</b>	234kD
<b>Cell Pathway</b>	[Isoform 1]: Cell membrane ; Single-pass type I membrane protein .; [Isoform 2]: Secreted .; [Isoform 3]: Secreted .
<b>Tissue Specificity</b>	Highly expressed in fetal kidney, and at slightly lower levels in fetal brain, lung and liver.
<b>Function</b>	disease:Overexpressed and constitutively tyrosine phosphorylated in colon, liver, pancreas and gastric carcinoma cell lines. Overexpression increases MET activation and promotes invasive growth.,function:Receptor for SEMA4D. Plays a role in RHOA activation and subsequent changes of the actin cytoskeleton. Plays a role in axon guidance, invasive growth and cell migration.,PTM:Phosphorylated on tyrosine residues by ERBB2 and MET upon SEMA4D binding.,PTM:Proteolytic processing favors heterodimerization with PLXNB2 and SEMA4D binding.,similarity:Belongs to the plexin family.,similarity:Contains 1 Sema domain.,similarity:Contains 3 IPT/TIG domains.,subunit:Monomer, and heterodimer with PLXNB2 after proteolytic processing. Binds RAC1 that has been activated by GTP binding. Interaction with SEMA4D promotes binding of

Nanjing BYabscience technology Co.,Ltd



cytoplasmic ligands. Binds PLXNA1 (By similarity). Binds ARHGEF11, ARHGEF12, E

**Background**

disease:Overexpressed and constitutively tyrosine phosphorylated in colon, liver, pancreas and gastric carcinoma cell lines. Overexpression increases MET activation and promotes invasive growth.,function:Receptor for SEMA4D. Plays a role in RHOA activation and subsequent changes of the actin cytoskeleton. Plays a role in axon guidance, invasive growth and cell migration.,PTM:Phosphorylated on tyrosine residues by ERBB2 and MET upon SEMA4D binding.,PTM:Proteolytic processing favors heterodimerization with PLXNB2 and SEMA4D binding.,similarity:Belongs to the plexin family.,similarity:Contains 1 Sema domain.,similarity:Contains 3 IPT/TIG domains.,subunit:Monomer, and heterodimer with PLXNB2 after proteolytic processing. Binds RAC1 that has been activated by GTP binding. Interaction with SEMA4D promotes binding of cytoplasmic ligands. Binds PLXNA1 (By similarity). Binds ARHGEF11, ARHGEF12, ERBB2, MET, MST1R, RND1, NRP1 and NRP2.,tissue specificity:Highly expressed in fetal kidney, and at slightly lower levels in fetal brain, lung and liver.,

**matters needing attention**

Avoid repeated freezing and thawing!

**Usage suggestions**

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

**Products Images**